

INNOVA JUNIOR COLLEGE  
JC 2 PRELIMINARY EXAMINATION  
in preparation for General Certificate of Education Advanced Level  
**Higher 2**

CANDIDATE  
NAME

CLASS

INDEX NUMBER

---

**CHEMISTRY**

**9729/01**

Paper 1 Multiple Choice

**14 Sep 2018**

**1 hour**

Additional Materials:

*Data Booklet*

Multiple Choice Answer Sheet

---

**READ THESE INSTRUCTIONS FIRST**

Write your index number, name and class on all the work you hand in.

Write in soft pencil.

Do not use staples, paper clips, highlighters, glue or correction fluid.

There are **thirty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

**Read the instructions on the Answer Sheet very carefully.**

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

---

This document consists of **16** printed pages and **1** blank page.



Innova Junior College

**[Turn over**

For each question there are four possible answers, **A**, **B**, **C**, and **D**. Choose the **one** you consider to be correct.

- 1 In an experiment, 0.002 mol acidified iron (II) sulfate is quantitatively oxidised by 0.001 mol of an oxidising agent,  $\text{ZO}_3^-$ . Given that  $\text{Fe}^{2+}$  is oxidised to  $\text{Fe}^{3+}$ , what is the final oxidation state of **Z**?
- A +1                      B +2                      C +3                      D +4
- 2 Gaseous particle **X** has a proton number  $n$  and a charge of +1.  
Gaseous particle **Y** has a proton number of  $(n + 1)$  and is isoelectronic with **X**.  
Which of the following statements correctly describe **X** and **Y**?
- 1 **Y** has a smaller radius than **X**.  
2 **Y** requires less energy than **X** when a further electron is removed from each particle.  
3 **Y** releases less energy than **X** when an electron is added to each particle.
- A 1 only  
B 2 only  
C 1 and 2 only  
D 2 and 3 only
- 3 What is the electronic configuration of an element with a **second** ionisation energy higher than that of each of the elements either side of it in the Periodic Table?
- A  $1s^2 2s^2 2p^6 3s^2$   
B  $1s^2 2s^2 2p^6 3s^2 3p^1$   
C  $1s^2 2s^2 2p^6 3s^2 3p^2$   
D  $1s^2 2s^2 2p^6 3s^2 3p^3$

- 4 The table shows the charge and radius of each of six ions.

ion	J <sup>+</sup>	L <sup>+</sup>	M <sup>2+</sup>	X <sup>-</sup>	Y <sup>-</sup>	Z <sup>2-</sup>
radius / nm	0.14	0.18	0.15	0.14	0.18	0.15


The ionic solids JX, LY and MZ are of the same lattice type. What is the correct order of their lattice energies, placing the most exothermic first?

- A** JX, MZ, LY  
**B** LY, MZ, JX  
**C** MZ, JX, LY  
**D** MZ, LY, JX
- 5 In certain microwave ovens, the wave energy produced is absorbed by certain polar molecules.

Which of the following would absorb microwave energy?

- 1 CH<sub>3</sub>CH<sub>2</sub>OH  
2 AlCl<sub>3</sub>  
3 CO<sub>2</sub>  
4 CH<sub>3</sub>F
- A** 1 and 2 only  
**B** 1 and 4 only  
**C** 2 and 3 only  
**D** 3 and 4 only


6




Which one of the following statements is true for the observation?

- A Gas E could be He while gas F could be  $\text{CO}_2$ .
- B Gas E could be  $\text{NH}_3$  while gas F could be  $\text{H}_2$ .
- C Gas E could form intermolecular hydrogen bonds while gas F could form intermolecular permanent dipole-permanent dipole.
- D Gas E could form stronger intermolecular instantaneous dipole-induced dipole than gas F.

- 7 The enzyme alcohol dehydrogenase catalyses an important step in the production of ethanol by fermentation.



The graph shows how the rate of this enzyme-catalysed reaction varies with the concentration of ethanal.




Which statement best explains the reason for the flattening off of the curve?

- A All the ethanal has been used up and the reaction has finished.
- B As the ethanol product builds up the reaction slows down.
- C At high ethanal concentrations all the active sites in the enzyme molecules are occupied by ethanal molecules.
- D At high ethanal concentrations the ethanal inhibits the action of enzyme.


- 8 The kinetics of the zero order reaction  $P \rightarrow Q$  were investigated under different conditions. The table shows pairs of quantities that were plotted as graphs.

Which pairs gave the following graph?



	y-axis	x-axis
1	rate	time
2	rate constant	time
3	rate constant	temperature
4	rate	concentration


- A 1 and 4 only  
 B 2 and 3 only  
 C 1, 2 and 3 only  
 D 1, 2 and 4 only
- 9 The graph below shows how the percentage of product present at equilibrium varies with temperature and pressure for a reaction.



Which reaction could the graph represent?

- A  $4\text{Fe(s)} + 3\text{O}_2\text{(g)} \rightleftharpoons 2\text{Fe}_2\text{O}_3\text{(s)}$   $\Delta H = -1644 \text{ kJ mol}^{-1}$   
 B  $2\text{C(s)} + \text{O}_2\text{(g)} \rightleftharpoons 2\text{CO(g)}$   $\Delta H = -222 \text{ kJ mol}^{-1}$   
 C  $\text{N}_2\text{O}_4\text{(g)} \rightleftharpoons 2\text{NO}_2\text{(g)}$   $\Delta H = +57.2 \text{ kJ mol}^{-1}$   
 D  $\text{CO(g)} + \text{Cl}_2\text{(g)} \rightleftharpoons \text{COCl}_2\text{(s)}$   $\Delta H = +86 \text{ kJ mol}^{-1}$

- 10 The standard enthalpy change of the following reaction is  $-896.4 \text{ kJ mol}^{-1}$ .




	NO(g)	H <sub>2</sub> O <sub>2</sub> (l)
$\Delta H_f^\theta / \text{kJ mol}^{-1}$	+ 90.3	- 187.8

Using the standard enthalpy change of formation in the above table, what is the standard enthalpy change of formation, in  $\text{kJ mol}^{-1}$ , of  $\text{HN}_3(\text{l})$ ?

- A +264  
 B +528  
 C +618  
 D +632
- 11 A student calculated the lattice energy for magnesium oxide using the cycle shown.

The enthalpy change of atomisation of magnesium,  $\Delta H_{\text{atomisation}}^\theta \text{Mg}$ , is the energy needed when 1 mol of gaseous magnesium atoms is formed from 1 mol of solid magnesium under standard conditions.



However, the value calculated by the student for the lattice energy was **less** exothermic than the correct value.

Which errors could have been made in the calculation?

- 1 omitting the enthalpy change of atomisation of magnesium
  - 2 omitting the first ionisation energy of magnesium
  - 3 using the standard enthalpy change of combustion of magnesium rather than the standard enthalpy change of formation of magnesium oxide
- A 1 and 2 only  
 B 1 and 3 only  
 C 2 and 3 only  
 D 1, 2 and 3

12 Which changes are accompanied by an increase in entropy of the system?

- 1 freezing of water
- 2 sublimation of iodine
- 3 vaporisation of ethanol
- 4 conversion of  $O_2(g)$  to  $O_3(g)$

- A 1 and 2 only  
B 2 and 3 only  
C 3 and 4 only  
D 1 and 4 only

13 The value of the ionic product of water,  $K_w$ , varies with temperature.

Temperature / °C	$K_w / \text{mol}^2 \text{dm}^{-6}$
25	$1.0 \times 10^{-14}$
62	$1.0 \times 10^{-13}$


What can be deduced from this information?

- A The ionic dissociation of water is an exothermic process.  
B The association of water molecules by hydrogen bonding increases as temperature increases.  
C The pH of pure water increases with temperature.  
D At 62 °C, water with a pH of 6.5 is neutral.



- 14 The following graph shows the changes in pH of 20.0 cm<sup>3</sup> of 1.0 mol dm<sup>-3</sup> acid solution when excess 1.0 mol dm<sup>-3</sup> alkali solution is added gradually.


Which one of the following pairs of solution with a suitable indicator could have resulted in the graph below?



	alkali	acid	indicator
<b>A</b>	NaOH	H <sub>2</sub> SO <sub>4</sub>	Methyl orange
<b>B</b>	NH <sub>3</sub>	HNO <sub>3</sub>	Phenolphthalein
<b>C</b>	KOH	CH <sub>3</sub> CO <sub>2</sub> H	Methyl orange
<b>D</b>	Ba(OH) <sub>2</sub>	CH <sub>3</sub> CH <sub>2</sub> CO <sub>2</sub> H	Phenolphthalein

- 15 Use of the Data Booklet is relevant to this question.

The AgCl / Ag reference electrode is based on the following reaction:




The reduction potential of the Fe<sup>2+</sup> / Fe half-cell was found to be -0.64 V when measured using the AgCl / Ag reference electrode.

What is the reduction potential of the AgCl / Ag electrode with respect to the standard hydrogen electrode?

- A** -1.10 V      **B** -0.20 V      **C** +0.20 V      **D** +1.10 V

- 16 Use of the Data Booklet is relevant to this question.


A cell is set up by connecting a  $\text{Cu}^{2+}/\text{Cu}$  half-cell and an acidified  $\text{MnO}_4^-/\text{Mn}^{2+}$  half-cell.



Which change in the set up will cause the corresponding effect on the e.m.f (potential difference) of the cell?

	change	effect on e.m.f of cell
A	addition of excess $\text{NH}_3$ (aq) into oxidation half-cell	increases
B	addition of concentrated $\text{H}_2\text{SO}_4$ into reduction half-cell	decreases
C	increase the size of the copper electrode	increases
D	replace copper with an alloy of copper and zinc	remains the same

- 17 In an experiment, a cell was set up to obtain pure copper from a copper-silver alloy as shown below. Electrode **V** is pure copper and electrode **U** is the copper-silver alloy.







When a current of 40.0 A flows through the electrolyte for 26.8 minutes, the mass of the anode changes by 26.47 g.

What is the percentage of silver by mass in the electrode **U**?




- A 20%
- B 40%
- C 60%
- D 80%

18 Which of the following is a planar molecule?







19 A hydrocarbon, on heating with an excess of hot concentrated acidic  $\text{KMnO}_4$  produces  $\text{HO}_2\text{CCH}_2\text{CH}_2\text{CH}_2\text{COCH}_3$ .

What could the hydrocarbon be?



- A 1 and 2 only  
 B 1 and 3 only  
 C 2 and 3 only  
 D 1, 2 and 3

- 20 When propene reacts with  $\text{Br}_2$  in the presence of excess aqueous  $\text{KNO}_3$ , what are the two major products formed?



- 21 The molecule of benzene,  $\text{C}_6\text{H}_6$  is a regular hexagon in which the  $\pi$  electrons are described as delocalised.




Which of the following statements are true for benzene?

- 1 It prevents benzene from undergoing addition reactions.
  - 2 All C–C bonds lengths in benzene are intermediate between C–C bond in an alkane and C=C bond in an alkene.
  - 3 The enthalpy change of hydrogenation of benzene is less exothermic than that predicted for cyclohexatriene.
  - 4 The presence of  $\pi$  electrons enables benzene to be a good electrical conductor.
- A 1 and 2 only  
 B 2 and 3 only  
 C 1, 2 and 3 only  
 D 1, 2 and 4 only




- 22 Some chlorobutanes were separately treated with hot ethanolic sodium hydroxide. Two of these gave the same hydrocarbon,  $C_4H_6$ .

From which pair of chlorobutanes was this hydrocarbon obtained?


- A  $CH_3CH_2CH_2CH_2Cl$  and  $CH_3CH_2CH_2CHCl_2$   
 B  $CH_3CH_2CH_2CH_2Cl$  and  $ClCH_2CH_2CH_2CH_2Cl$   
 C  $CH_3CH_2CCl_2CH_3$  and  $ClCH_2CH_2CH_2CHCl_2$   
 D  $CH_3CHClCHClCH_3$  and  $ClCH_2CH_2CH_2CH_2Cl$
- 23 Deuterium, D, is the  $^2_1H$  isotope of hydrogen.  
 Which of the following reactions yield a carbon compound containing deuterium?



- A 1 and 2 only  
 B 1 and 4 only  
 C 1, 2 and 3 only  
 D 2, 3 and 4 only
- 24 Which of the following synthetic routes **does not** produce ethanoic acid?





- 25 Hydrocortisone is a steroid hormone produced by the adrenal gland and is released in response to stress. It is commonly used as an active ingredient in anti-inflammatory creams.



Hydrocortisone

Which of the following statements about Hydrocortisone is true?


- A** When treated with  $\text{NaBH}_4$  in the presence of methanol, it forms a compound containing 5 hydroxy groups.
- B** When warmed with aqueous alkaline iodine, a yellow precipitate is observed.
- C** When treated with cold dilute  $\text{KMnO}_4$ , it forms a compound containing 2 hydroxy groups.
- D** When treated with an excess of hot concentrated acidified  $\text{KMnO}_4$ , it forms a compound containing 7 chiral centres.
- 26 Equal amounts of compounds **P**, **Q**, **R** and **S** are separately shaken with  $100 \text{ cm}^3$  of water. The pH of each resultant solution is then measured.

**P****Q****R****S**

Which of the following shows the correct order of increasing order of acidity of the solutions formed?

- A** P, R, S, Q
- B** P, S, R, Q
- C** Q, P, S, R
- D** Q, S, R, P


27 The diagram shows the structure of the tetrapeptide **T**.



When 0.1 mole of **T** is heated under reflux with  $\text{NaOH}(\text{aq})$  until no further reaction occurs, how many moles of  $\text{NaOH}$  will react?

- A 0.4  
B 0.5  
C 0.6  
D 0.7

28 Lysine is an  $\alpha$ -amino acid.



Which structure is predominant when lysine is in an aqueous solution of pH 9.5, given that lysine has three  $\text{p}K_a$  values of 2.2, 8.9 and 10.5?

- |          |  |          |   |
|----------|--|----------|---|
| <b>A</b> | $\begin{array}{c} \text{H}_3\text{N}^+\text{CHCO}_2\text{H} \\   \\ (\text{CH}_2)_4 \\   \\ \text{NH}_3^+ \end{array}$ | <b>B</b> | $\begin{array}{c} \text{H}_2\text{NCHCO}_2^- \\   \\ (\text{CH}_2)_4 \\   \\ \text{NH}_2 \end{array}$   |
| <b>C</b> | $\begin{array}{c} \text{H}_3\text{N}^+\text{CHCO}_2^- \\   \\ (\text{CH}_2)_4 \\   \\ \text{NH}_3^+ \end{array}$       | <b>D</b> | $\begin{array}{c} \text{H}_2\text{NCHCO}_2^- \\   \\ (\text{CH}_2)_4 \\   \\ \text{NH}_3^+ \end{array}$ |

**29** Element **J** is in Period 3 of the Periodic Table. The following three statements are the properties of element **J** or its compounds.

- Adding  $\text{NaOH(aq)}$  to the solution resulting from the reaction of a chloride of **J** with water produces a white precipitate which is soluble in an excess of  $\text{NaOH(aq)}$ .
- Element **J** is a solid at room temperature.
- The oxide of element **J** is soluble in hydrochloric acid.

What is the identity of element **J**?

- |                    |                     |
|--------------------|---------------------|
| <b>A</b> magnesium | <b>C</b> silicon    |
| <b>B</b> aluminium | <b>D</b> phosphorus |

**30** The use of the Data Booklet is relevant to this question.

The colours of various vanadium ions in aqueous solution are given in the table below.

Oxidation state	V	IV	III	II
Ion	$\text{VO}_2^+$	$\text{VO}^{2+}$	$\text{V}^{3+}$	$\text{V}^{2+}$
Colour	yellow	blue	green	violet


What is the final colour of the solution when excess zinc powder is added to an acidified solution containing  $\text{VO}_2^+$  ions?

- |                 |                 |
|-----------------|-----------------|
| <b>A</b> yellow | <b>C</b> green  |
| <b>B</b> blue   | <b>D</b> violet |

**End of Paper**







INNOVA JUNIOR COLLEGE  
JC 2 PRELIMINARY EXAMINATION  
in preparation for General Certificate of Education Advanced Level  
**Higher 2**

CANDIDATE  
NAME

**MARK SCHEME**

CLASS

INDEX NUMBER

**CHEMISTRY**

**9729/01**

Paper 1 Multiple Choice

**XX XXX 2018**

**1 hour**

Additional Materials:

*Data Booklet*

Multiple Choice Answer Sheet

**READ THESE INSTRUCTIONS FIRST**

Write your index number, name and class on all the work you hand in.

Write in soft pencil.

Do not use staples, paper clips, highlighters, glue or correction fluid.

There are **thirty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

**Read the instructions on the Answer Sheet very carefully.**

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

1	<b>C</b>	6	<b>A</b>	11	<b>A</b>	16	<b>A</b>	21	<b>C</b>	26	<b>B</b>
2	<b>A</b>	7	<b>C</b>	12	<b>B</b>	17	<b>A</b>	22	<b>D</b>	27	<b>D</b>
3	<b>B</b>	8	<b>D</b>	13	<b>D</b>	18	<b>D</b>	23	<b>C</b>	28	<b>D</b>
4	<b>C</b>	9	<b>B</b>	14	<b>D</b>	19	<b>C</b>	24	<b>D</b>	29	<b>B</b>
5	<b>B</b>	10	<b>A</b>	15	<b>C</b>	20	<b>D</b>	25	<b>A</b>	30	<b>D</b>

A: 7 B: 6 C:7 D: 10

This document consists of **X** printed pages.



Innova Junior College

**[Turn over**

For each question there are four possible answers, **A**, **B**, **C**, and **D**. Choose the **one** you consider to be correct.

- 1 In an experiment, 0.002 mol acidified iron (II) sulfate is quantitatively oxidised by 0.001 mol of an oxidising agent,  $\text{ZO}_3^-$ . Given that  $\text{Fe}^{2+}$  is oxidised to  $\text{Fe}^{3+}$ , what is the final oxidation state of **Z**?

**A** +1                      **B** +2                      **C** +3                      **D** +4

Answer: C

Amount of electrons released by  $\text{Fe}^{2+}$  = 0.002 mol  
= amount of electrons accepted by  $\text{YO}_3^-$

Amount of  $\text{YO}_3^-$ ; amount of e = 1 : 2

Oxidation number of Y in  $\text{YO}_3^-$  = +5

New oxidation number of Y = +5 – 2 = +3

- 2 Gaseous particle **X** has a proton number  $n$  and a charge of +1.

Gaseous particle **Y** has a proton number of  $(n + 1)$  and is isoelectronic with **X**.

Which of the following statements correctly describe **X** and **Y**?

- 1** Y has a smaller radius than **X**.  
**2** Y requires less energy than **X** when a further electron is removed from each particle.  
**3** Y releases less energy than **X** when an electron is added to each particle.

- A** 1 only  
**B** 2 only  
**C** 1 and 2 only  
**D** 2 and 3 only

Answer: A

particle	proton number	electron number	charge
<b>X</b>	$n$	$n-1$	+1
<b>Y</b>	$n+1$	$n-1$	+2

Since **Y** has higher nuclear charge than **X**, and both **X** and **Y** have same shielding effect (same electron number), **Y** has higher effective nuclear charge and valence electrons are more strongly attracted to the nucleus.

Thus, **Y** has smaller radius than **X**. (Option 1 is correct)

**Y** requires more energy than **X** when a further electron is removed from each particle. (Option 2 is wrong)

**Y** releases more energy than **X** when an electron is added to each particle. (Option 3 is wrong)

- 3 What is the electronic configuration of an element with a **second** ionisation energy higher than that of each of the elements either side of it in the Periodic Table?

- A  $1s^22s^22p^63s^2$   
**B**  $1s^22s^22p^63s^23p^1$   
 C  $1s^22s^22p^63s^23p^2$   
 D  $1s^22s^22p^63s^23p^3$

Answer: B

The electronic configuration for second ionisation energy of the above species:

- A:  $1s^22s^22p^63s^1$   
 B:  $1s^22s^22p^63s^2$   
 C:  $1s^22s^22p^63s^23p^1$   
 D:  $1s^22s^22p^63s^23p^2$

As there is a decrease in ionisation energy between the species between  $ns^2$  and  $ns^2np^1$ , species that have electronic configuration of  $ns^2$  will have second ionisation energy higher than that of each of the elements either side of it ( $ns^1$  and  $ns^2np^1$ ) in the Periodic Table.

- 4 The table shows the charge and radius of each of six ions.

ion	J <sup>+</sup>	L <sup>+</sup>	M <sup>2+</sup>	X <sup>-</sup>	Y <sup>-</sup>	Z <sup>2-</sup>
radius / nm	0.14	0.18	0.15	0.14	0.18	0.15

The ionic solids JX, LY and MZ are of the same lattice type. What is the correct order of their lattice energies, placing the most exothermic first?

- A JX, MZ, LY  
 B LY, MZ, JX  
**C** MZ, JX, LY  
 D MZ, LY, JX

Answer: C

Since  $\left| \text{Lattice energy, } \Delta H_{\text{latt}}^\theta \right| \propto \left| \frac{q_+ q_-}{r_+ + r_-} \right|$ , compound MZ has the most exothermic lattice

energy as M<sup>2+</sup> and Z<sup>2-</sup> has higher charge than that of J<sup>+</sup>, L<sup>+</sup>, X<sup>-</sup> and Y<sup>-</sup>. Compound LY has the least exothermic lattice energy as L<sup>+</sup> and Y<sup>-</sup> has larger ionic radius than that of J<sup>+</sup> and Y<sup>-</sup>.

- 5 In certain microwave ovens, the wave energy produced is absorbed by certain polar molecules.

Which of the following would absorb microwave energy?

- 1  $\text{CH}_3\text{CH}_2\text{OH}$
  - 2  $\text{AlCl}_3$
  - 3  $\text{CO}_2$
  - 4  $\text{CH}_3\text{F}$
- A 1 and 2 only
- B** 1 and 4 only
- C 2 and 3 only
- D 3 and 4 only


Answer: B

Option 2 is non polar molecule.  $\text{AlCl}_3$  is trigonal planar in shape and there is no net dipole moment in the molecule as all the dipole moments cancel out each other.

Option 3 is non polar molecule.  $\text{CO}_2$  is linear in shape and there is no net dipole moment in the molecule as all the dipole moments cancel out each other.

Option 1 and 4 are polar as there is net dipole moment in the molecule.

6



Which one of the following statements is true for the observation?

- A** Gas **E** could be He while gas **F** could be  $\text{CO}_2$ .
- B** Gas **E** could be  $\text{NH}_3$  while gas **F** could be  $\text{H}_2$ .
- C** Gas **E** could form intermolecular hydrogen bonds while gas **F** could form intermolecular permanent dipole-permanent dipole.
- D** Gas **E** could form stronger intermolecular instantaneous dipole-induced dipole than gas **F**.

Answer: A

$$pV = nRT$$

$$1.5 \times 10^{13} 25 \times V = 1 \times 8.31 \times (273+27)$$

$$V = 0.0164 \text{ m}^3$$

$$V = 16.4 \text{ dm}^3$$

From the above calculation, it can be deduced that gas **E** behave more ideally than gas **F**.


Option A is correct. He behaves more like an ideal gas as it has smaller electron cloud size and weaker i.d.-i.d. than that of  $\text{CO}_2$ . Intermolecular forces of attraction between He is negligible.

Option B is wrong.  $\text{NH}_3$  can form hydrogen bonds between molecules and there is significant intermolecular forces of attraction.  $\text{NH}_3$  deviates more from ideal gas.

Option C is wrong. Gas **E** should form weaker i.d.-i.d. between molecules than that of **F** as gas **E** behave more ideally than gas **F**.

Option D is wrong. Gas **E** should form weaker i.d.-i.d. than gas **F** as gas **E** behave more ideally than gas **F**.

- 7 The enzyme alcohol dehydrogenase catalyses an important step in the production of ethanol by fermentation.



The graph shows how the rate of this enzyme-catalysed reaction varies with the concentration of ethanal.

[ethanal]

Which statement best explains the reason for the flattening off of the curve?

- A All the ethanal has been used up and the reaction has finished.
- B As the ethanol product builds up the reaction slows down.
- C** At high ethanal concentrations all the active sites in the enzyme molecules are occupied by ethanal molecules.
- D At high ethanal concentrations the ethanal inhibits the action of enzyme.

Answer: C


Option A is wrong as there is an increase in the concentration of ethanal. The ethanal has not been used up and the reaction has not finished.

Option B is wrong as the concentration of the ethanol product builds up the reaction will not slow down the rate of reaction.

Option D is wrong as ethanal is not an inhibitor but substrate (reactant) of this reaction.

- 8 The kinetics of the zero order reaction  $P \rightarrow Q$  were investigated under different conditions. The table shows pairs of quantities that were plotted as graphs.

Which pairs gave the following graph?



	y-axis	x-axis
1	rate	time
2	rate constant	time
3	rate constant	temperature
4	rate	concentration

- A** 1 and 4 only  
**B** 2 and 3 only  
**C** 1, 2 and 3 only  
**D** 1, 2 and 4 only


Answer: D

For a zero order reaction, rate of the reaction is independent of time and concentration.

Also, rate constant is dependent on temperature and activation energy (from Arrhenius equation.)



- 9 The graph below shows how the percentage of product present at equilibrium varies with temperature and pressure for a reaction.



Which reaction could the graph represent?

- A  $4\text{Fe (s)} + 3\text{O}_2\text{(g)} \rightleftharpoons 2\text{Fe}_2\text{O}_3\text{(s)}$   $\Delta H = -1644 \text{ kJ mol}^{-1}$
- B**  $2\text{C (s)} + \text{O}_2\text{(g)} \rightleftharpoons 2\text{CO (g)}$   $\Delta H = -222 \text{ kJ mol}^{-1}$
- C  $\text{N}_2\text{O}_4\text{(g)} \rightleftharpoons 2\text{NO}_2\text{(g)}$   $\Delta H = +57.2 \text{ kJ mol}^{-1}$
- D  $\text{CO(g)} + \text{Cl}_2\text{(g)} \rightleftharpoons \text{COCl}_2\text{(s)}$   $\Delta H = +86 \text{ kJ mol}^{-1}$


Answer: B

From the graph, when the pressure increases, the proportion of product molecules decrease. This indicates that when pressure increases, the equilibrium position shifts to left (which have fewer gaseous molecules to decrease the pressure). Option A and option D is wrong.

From the graph, when the temperature increases, the proportion of product molecules decrease too. This indicates that when temperature increases, the equilibrium position shifts to left (which is endothermic to reduce the temperature. Since option B is exothermic in the forward reaction, it would be endothermic in the backward reaction.

Only Option B shows the correct trend when temperature and pressure increase.

- 10 The standard enthalpy change of the following reaction is  $-896.4 \text{ kJ mol}^{-1}$ .



	NO(g)	H <sub>2</sub> O <sub>2</sub> (l)
$\Delta H_f^\ominus/\text{kJ mol}^{-1}$	+ 90.3	- 187.8

Using the standard enthalpy change of formation in the above table, what is the standard enthalpy change of formation, in  $\text{kJ mol}^{-1}$ , of  $\text{HN}_3(\text{l})$ ?

- A** +264  
**B** +528  
**C** +618  
**D** +632

Answer: A

$$\text{Using } \Delta H_r^\ominus = \sum n\Delta H_f^\ominus (\text{products}) - \sum m\Delta H_f^\ominus (\text{reactants})$$


$$\Delta H_r^\ominus = \Delta H_f (\text{H}_2\text{O}_2) - [2 \Delta H_f (\text{HN}_3) + 2 \Delta H_f (\text{NO})]$$

$$-896.4 = -187.8 - (2 \Delta H_f (\text{HN}_3) + 2 \times 90.3)$$

$$\Delta H_f (\text{HN}_3) = +264 \text{ kJ mol}^{-1}$$

- 11 A student calculated the lattice energy for magnesium oxide using the cycle shown.

The enthalpy change of atomisation of magnesium,  $\Delta H_{\text{atomisation}}^{\theta} \text{Mg}$ , is the energy needed when 1 mol of gaseous magnesium atoms is formed from 1 mol of solid magnesium under standard conditions.



However, the value calculated by the student for the lattice energy was **less** exothermic than the correct value.

Which errors could have been made in the calculation?

- 1 omitting the enthalpy change of atomisation of magnesium
- 2 omitting the first ionisation energy of magnesium
- 3 using the standard enthalpy change of combustion of magnesium rather than the standard enthalpy change of formation of magnesium oxide

- A** 1 and 2 only  
**B** 1 and 3 only  
**C** 2 and 3 only  
**D** 1, 2 and 3

Ans: A

By Hess's law,

$$\text{Lattice energy} = - [\Delta H_{\text{atomisation}}^{\theta} \text{Mg}] - [1^{\text{st}} \text{ and } 2^{\text{nd}} \text{ I.E. of Mg}] - 950 \\ + \Delta H_{\text{formation}}^{\theta} \text{MgO}$$

1 is correct as the lattice energy will be less exothermic if the enthalpy change of atomisation of Mg is omitted. This is because, the enthalpy change of atomisation of Mg is a positive term.

2 is correct as the lattice energy will be less exothermic if the first ionisation energy of Mg is omitted. This because, the first ionisation energy of Mg is a positive term

3 is incorrect as the replacing the standard enthalpy change of formation of MgO with the standard enthalpy change of combustion of Mg will have no effect on the value of the lattice energy. This is because both values are the same.

12 Which changes are accompanied by an increase in entropy of the system?

- 1 freezing of water
- 2 sublimation of iodine
- 3 vaporisation of ethanol
- 4 conversion of  $O_2(g)$  to  $O_3(g)$

- A 1 and 2 only  
**B** 2 and 3 only  
 C 3 and 4 only  
 D 1 and 4 only

Answer: B

For option 2 and 3, there is an increase in the number of gaseous molecules (from solid to gas and from liquid to gas), there is an increase in the disorderness of the system, entropy increases.

For option 1, there is a decrease in the disorderness of the system (from liquid to solid) and similarly for that of option 4.  $3O_2(g) \rightarrow 2O_3(g)$

13 The value of the ionic product of water,  $K_w$ , varies with temperature.

Temperature / °C	$K_w / \text{mol}^2 \text{dm}^{-6}$
25	$1.0 \times 10^{-14}$
62	$1.0 \times 10^{-13}$

What can be deduced from this information?

- A The ionic dissociation of water is an exothermic process.  
 B The association of water molecules by hydrogen bonding increases as temperature increases.  
 C The pH of pure water increases with temperature.  
**D** At 62 °C, water with a pH of 6.5 is neutral.

Answer: D

Option A is wrong as ionic dissociation of water involves bond breaking of water molecules, this is an endothermic reaction.


Option B is wrong. As temperature increases, more hydrogen bonds between molecules are broken.

Option C is wrong. The pH of pure water decreases with temperature. (pH = 7 at 25 °C and pH = 6.5 at 62 °C)

Option D is correct as water is neutral at all temperature as  $[H^+] = [OH^-]$ .

- 14 The following graph shows the changes in pH of 20.0 cm<sup>3</sup> of 1.0 mol dm<sup>-3</sup> acid solution when excess 1.0 mol dm<sup>-3</sup> alkali solution is added gradually.

Which one of the following pairs of solution with a suitable indicator could have resulted in the graph below?




	alkali	acid	indicator
A	NaOH	H <sub>2</sub> SO <sub>4</sub>	Methyl orange
B	NH <sub>3</sub>	HNO <sub>3</sub>	Phenolphthalein
C	KOH	CH <sub>3</sub> CO <sub>2</sub> H	Methyl orange
D	Ba(OH) <sub>2</sub>	CH <sub>3</sub> CH <sub>2</sub> CO <sub>2</sub> H	Phenolphthalein

Answer: D

Since the volume of alkali needed for neutralisation is half of that of acid, this indicates that there should be 2 OH<sup>-</sup> per base molecules. Since this is a weak acid-strong base titration, pH at equivalence point should be greater than 7 as a basic salt is formed. Phenolphthalein is a suitable indicator as its working range is greater than 7.

- 15 Use of the Data Booklet is relevant to this question.

The AgCl / Ag reference electrode is based on the following reaction:



The reduction potential of the Fe<sup>2+</sup> / Fe half-cell was found to be -0.64 V when measured using the AgCl / Ag reference electrode.

What is the reduction potential of the AgCl / Ag electrode with respect to the standard hydrogen electrode?

- A -1.10 V      B -0.20 V      C +0.20 V      D +1.10 V

Answer: C

The reduction potential of the Fe<sup>2+</sup> / Fe half-cell was found to be -0.44 V when measured using the standard hydrogen electrode.


$$E_{\text{cell}}^{\circ} = E_{\text{red}}^{\circ} - E_{\text{ox}}^{\circ}$$

$$-0.44 \text{ V} = E_{\text{red}}^{\circ} - 0.64$$

$$E_{\text{red}}^{\circ} = + 0.20 \text{ V}$$

16 Use of the Data Booklet is relevant to this question.

A cell is set up by connecting a  $\text{Cu}^{2+}/\text{Cu}$  half-cell and an acidified  $\text{MnO}_4^-/\text{Mn}^{2+}$  half-cell.



Which change in the set up will cause the corresponding effect on the e.m.f (potential difference) of the cell?

	change	effect on e.m.f of cell
<b>A</b>	addition of excess $\text{NH}_3$ (aq) into oxidation half-cell	increases
<b>B</b>	addition of concentrated $\text{H}_2\text{SO}_4$ into reduction half-cell	decreases
<b>C</b>	increase the size of the copper electrode	increases
<b>D</b>	replace copper with an alloy of copper and zinc	remains the same

Answer: A



$$E^\circ_{\text{cell}} = E^\circ_{\text{red}} - E^\circ_{\text{ox}} = +1.52 - (+0.34) = +1.18 \text{ V}$$


Option A is correct: when excess  $\text{NH}_3$  (aq) into oxidation half-cell,  $[\text{Cu}^{2+}]$  decreases as  $\text{Cu}^{2+}$  reacts with  $\text{NH}_3$  to form  $[\text{Cu}(\text{NH}_3)_4]^{2+}$ . Since  $[\text{Cu}^{2+}]$  decreases, equilibrium position shifts to the left,  $E^\circ_{\text{ox}}$  is less positive,  $E^\circ_{\text{cell}}$  becomes more positive.

Option B is wrong. When concentrated  $\text{H}_2\text{SO}_4$  is added into reduction half-cell,  $[\text{H}^+]$  increases. Since  $[\text{H}^+]$  increases, equilibrium position shifts to the right,  $E^\circ_{\text{red}}$  is more positive,  $E^\circ_{\text{cell}}$  becomes more positive.

Option C is wrong. Increase the size of the copper electrode will not have effect on e.m.f. of the cell.

Option D is wrong. Replace copper with an alloy of copper and zinc will result in more positive  $E^\circ_{\text{cell}}$ .  $(+1.52 - (-0.76)) = +2.28 \text{ V}$

- 17 In an experiment, a cell was set up to obtain pure copper from a copper-silver alloy as shown below. Electrode **V** is pure copper and electrode **U** is the copper-silver alloy.



When a current of 40.0 A flows through the electrolyte for 26.8 minutes, the mass of the anode changes by 26.47 g.

What is the percentage of silver by mass in the electrode **U**?

- A** 20%
- B** 40%
- C** 60%
- D** 80%

Answer: A

$$I \times t = n_e \times F$$

$$40 \times 26.8 \times 60 = n_e \times 96500$$

$$n_e = 0.6665 \text{ mol}$$





Since  $\text{Cu} : e = 1 : 2$ ,

$$\text{Amount of Cu formed} = \frac{1}{2} \times 0.6665 = 0.3332 \text{ mol}$$

$$\text{Mass of Cu formed} = 0.3332 \times 63.5 = 21.16 \text{ g}$$

$$\% \text{ silver by mass} = (26.47 - 21.16) / 26.47 \times 100\% = 20\%$$

18 Which of the following is a planar molecule?



Answer: D

Option A is wrong. There are 2  $\text{sp}^3$  carbon atoms (tetrahedral around the C) in the molecule.

Option B is wrong. The molecule is trigonal pyramidal around the N atom.


Option C is wrong. There are 1  $\text{sp}^3$  carbon atom (tetrahedral around the C) in the molecule.

Option D is correct. All the carbons are  $\text{sp}^2$  carbon in the molecule.



- 19 A hydrocarbon, on heating with an excess of hot concentrated acidic  $\text{KMnO}_4$  produces  $\text{HO}_2\text{CCH}_2\text{CH}_2\text{CH}_2\text{COCH}_3$ .


What could the hydrocarbon be?




- A 1 and 2 only  
 B 1 and 3 only  
 C 2 and 3 only  
 D 1, 2 and 3

Answer: C


Option A is wrong.




Option B is correct.




Option C is correct.



- 20 When propene reacts with  $\text{Br}_2$  in the presence of excess aqueous  $\text{KNO}_3$ , what are the two major products formed?



Answer: D



- 21 The molecule of benzene,  $C_6H_6$  is a regular hexagon in which the  $\pi$  electrons are described as delocalised.

Which of the following statements are true for benzene?

- 1 It prevents benzene from undergoing addition reactions.
  - 2 All C–C bonds lengths in benzene are intermediate between C–C bond in an alkane and C=C bond in an alkene.
  - 3 The enthalpy change of hydrogenation of benzene is less exothermic than that predicted for cyclohexatriene.
  - 4 The presence of  $\pi$  electrons enables benzene to be a good electrical conductor.
- A 1 and 2 only  
B 2 and 3 only  
**C** 1, 2 and 3 only  
D 1, 2 and 4 only

Answer: C

Option 1 is correct. Benzene undergoes substitution reaction to remain its aromaticity.

Option 2 is correct. As the  $\pi$  electrons are delocalised in the benzene ring, all the C-C bonds have partial double bond character.

Option 3 is correct. As the  $\pi$  electrons are delocalised in the benzene ring, benzene is more stable and it occupies lower energy level as it is more stable due to its aromaticity. As a result, less energy is released from hydrogenation of benzene.

Option 4 is wrong. Although the  $\pi$  electrons are delocalised in the benzene ring, there is no free mobile electrons and ions in liquid benzene, thus benzene is not a good conductor of electricity.

- 22 Some chlorobutanes were separately treated with hot ethanolic sodium hydroxide. Two of these gave the same hydrocarbon,  $C_4H_6$ .



From which pair of chlorobutanes was this hydrocarbon obtained?

- A  $CH_3CH_2CH_2CH_2Cl$  and  $CH_3CH_2CH_2CHCl_2$   
 B  $CH_3CH_2CH_2CH_2Cl$  and  $ClCH_2CH_2CH_2CH_2Cl$   
 C  $CH_3CH_2CCl_2CH_3$  and  $ClCH_2CH_2CH_2CHCl_2$   
 D  $CH_3CHClCHClCH_3$  and  $ClCH_2CH_2CH_2CH_2Cl$

Answer: D




For option A, B and C, there is at least one molecule in the option contain 1 or 3 Cl atoms which after elimination can only form 1 or 3 C=C bond.

For option D, the two compounds can produce the same compound, as shown below:



23 Deuterium, D, is the  ${}^2_1\text{H}$  isotope of hydrogen.

Which of the following reactions yield a carbon compound containing deuterium?



A 1 and 2 only


B 1 and 4 only

**C** 1, 2 and 3 only


D 2, 3 and 4 only

Answer: C


Option 1 is correct.




Option 2 is correct.







Option 3 is correct.



Option 4 is wrong.




24 Which of the following synthetic routes **does not** produce ethanoic acid?




Answer: D


Option A is correct.




Option B is correct.




Option C is correct.



Option D is wrong.



- 25 Hydrocortisone is a steroid hormone produced by the adrenal gland and is released in response to stress. It is commonly used as an active ingredient in anti-inflammatory creams.



Hydrocortisone

Which of the following statements about hydrocortisone is true?

- A** When treated with  $\text{NaBH}_4$  in the presence of methanol, it forms a compound containing 5 hydroxy groups.
- B** When warmed with aqueous alkaline iodine, a yellow precipitate is observed.
- C** When treated with cold dilute  $\text{KMnO}_4$ , it forms a compound containing 2 hydroxy groups.
- D** When treated with an excess of hot concentrated acidified  $\text{KMnO}_4$ , it forms a compound containing 7 chiral centres.


Answer: A

Option A is correct. The two ketone is reduced to secondary alcohol. Together with the other three alcohol in the molecule, the molecule has 5 hydroxy group.



Option B is wrong. There is no  $\text{H} / \text{R} - \overset{\text{O}}{\parallel} \text{C} - \text{CH}_3$  or  $\begin{array}{c} \text{H} \\ | \\ \text{R} / \text{H} - \text{C} - \text{OH} \\ | \\ \text{CH}_3 \end{array}$  in the molecule.

Option C is wrong. When treated with cold dilute  $\text{KMnO}_4$ , the alkene in the molecule will be converted to diol. Together with the other three alcohol in the molecule, the molecule has 5 hydroxy group.

Option D is wrong. When treated with an excess of hot concentrated acidified  $\text{KMnO}_4$ , it forms a compound containing 6 chiral centres.




- 26 Equal amounts of compounds **P**, **Q**, **R** and **S** are separately shaken with 100 cm<sup>3</sup> of water. The pH of each resultant solution is then measured.

**P****Q****R****S**

Which of the following shows the correct order of increasing order of acidity of the solutions formed?

- A** P, R, S, Q  
**B** P, S, R, Q  
**C** Q, P, S, R  
**D** Q, S, R, P

Answer: B




Solution is most acidic.

Cl group is more electronegative than Br and the negative charge of  $\text{C}/\text{CH}_2\text{CH}_2\text{CO}_2^-$  is dispersed to a greater extent than  $\text{BrCH}_2\text{CH}_2\text{CO}_2^-$ . Thus,  $\text{C}/\text{CH}_2\text{CH}_2\text{CO}_2^-$  is more stable than  $\text{BrCH}_2\text{CH}_2\text{CO}_2^-$ .  $\text{C}/\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$  is more acidic than  $\text{BrCH}_2\text{CH}_2\text{CO}_2\text{H}$ .

$\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$  is the least acidic among the four compounds.



27 The diagram shows the structure of the tetrapeptide **T**.



When 0.1 mole of **T** is heated under reflux with NaOH(aq) until no further reaction occurs, how many moles of NaOH will react?


**A** 0.4

**B** 0.5

**C** 0.6


**D** 0.7

Answer: D




The circled functional group can react with NaOH(aq), heat with reflux.


28 Lysine is an  $\alpha$ -amino acid.




Which structure is predominant when lysine is in an aqueous solution of pH 9.5, given that lysine has three  $pK_a$  values of 2.2, 8.9 and 10.5?



Answer: D



When  $\text{pH} > pK_a$ , the functional group circled will be deprotonated.  
When  $\text{pH} < pK_a$ , the functional group squared will be protonated.



29 Element J is in Period 3 of the Periodic Table. The following three statements are the properties of element J or its compounds.


- Adding NaOH(aq) to the solution resulting from the reaction of a chloride of J with water produces a white precipitate which is soluble in an excess of NaOH(aq).
- Element J is a solid at room temperature.
- The oxide of element J is soluble in hydrochloric acid.

What is the identity of element J?

- |   |           |   |            |
|---|-----------|---|------------|
| A | magnesium | C | silicon    |
| B | aluminium | D | phosphorus |

Answer: B

Based on the information provided, the only possible Period 3 element for J is Al.



Al is a solid at room temperature.

The oxide of Al is soluble in hydrochloric acid.



30 The use of the Data Booklet is relevant to this question.


The colours of various vanadium ions in aqueous solution are given in the table below.

Oxidation state	V	IV	III	II
Ion	$\text{VO}_2^+$	$\text{VO}^{2+}$	$\text{V}^{3+}$	$\text{V}^{2+}$
Colour	yellow	blue	green	violet

What is the final colour of the solution when excess zinc powder is added to an acidified solution containing  $\text{VO}_2^+$  ions?

- |   |        |   |        |
|---|--------|---|--------|
| A | yellow | C | green  |
| B | blue   | D | violet |


Answer: D



$$E^\circ_{\text{cell}} = E^\circ_{\text{red}} - E^\circ_{\text{ox}} = E^\circ_{\text{red}} - (-0.76)$$

For reaction to be spontaneous,  $E^\circ_{\text{cell}} > 0$  V. In order for  $E^\circ_{\text{cell}}$  to be  $> 0$ ,  $E^\circ_{\text{red}}$  must be  $> -0.76$ . Thus,  $\text{VO}_2^+$  will be eventually reduced to  $\text{V}^{2+}$ .

End of Paper



INNOVA JUNIOR COLLEGE  
 JC2 PRELIMINARY EXAMINATION  
 in preparation for General Certificate of Education Advanced Level  
**Higher 2**

CANDIDATE NAME

CLASS

INDEX NUMBER

## CHEMISTRY

**9729/02**

Paper 2 Structured Questions

**27 August 2018**

Candidates answer on the question paper.

**2 hours**

Additional Materials: *Data Booklet*

### READ THESE INSTRUCTIONS FIRST

Write your index number, name and civics group on all the work you hand in.  
 Write in dark blue or black pen.  
 You may use pencil for any diagrams, graphs or rough working.  
 Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions in the space provided.

You are advised to show all working in calculations.  
 You are reminded of the need for good English and clear presentation in your answers.  
 You are reminded of the need for good handwriting.  
 Your final answers should be in 3 significant figures.

You may use a calculator.


The number of marks is given in brackets [ ] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

For Examiner's Use	
1	9
2	8
3	6
4	15
5	13
6	12
7	12
<b>Significant Figures and Units</b>	
<b>Handwriting and Presentation</b>	
<b>Total</b>	<b>75</b>

This document consists of **19** printed pages **1** blank page.

- 1 (a) The following diagram shows the second ionisation energy of some elements.



- (i) Explain why the second ionisation energies of elements O, S, Se and Te show a decreasing trend.

.....  
 .....  
 .....[1]

- (ii) Explain why the second ionisation energy of Br is lower than that of Se.


.....  
 .....[1]

- (iii) Write the full ground state electronic configuration for Cr.

.....[1]

(iv) Draw the  $d_{xy}$  orbital in the axes provided below.

[1]



(b) Oxygen–oxygen bond lengths in some molecules are given below:

Molecule	Bond Length
Oxygen, $O_2$	0.121 nm
Hydrogen peroxide, $H_2O_2$	0.149 nm
Ozone, $O_3$	0.128 nm

(i) Draw the structure of the molecule ozone,  $O_3$ .

[1]

(ii) With reference to the data given above, comment and explain the oxygen–oxygen bond length in ozone as compared to oxygen and hydrogen peroxide.

.....

.....

.....

.....[2]

- (iii) Explain clearly why hydrogen peroxide is a liquid while oxygen is a gas at room conditions in terms of structure and bonding.

.....

.....


.....

.....[2]


[Total: 9]

- 2 Compounds containing nitrogen are important to life and have applications in science and medicine.


Three nitrogenous bases have the following structures.



*N*,4-dimethylbenzenamine



4-methylphenylamine



benzylamine

- (a) (i) Arrange the three bases above in increasing order of  $pK_b$ .

.....[1]

- (ii) Explain in terms of their molecular structures why benzylamine and 4-methylphenylamine have different  $pK_b$  values.

.....

.....

.....

.....

.....[2]

- (iii) Outline how *N*,4-dimethylbenzenamine may be produced from 4-methylphenylamine.

.....  
.....[1]

- (b) The  $pK_b$  of benzylamine is 4.66.

25.0 cm<sup>3</sup> of 0.025 mol dm<sup>-3</sup> benzylamine was completely neutralised by dilute hydrochloric acid of the same concentration. The salt formed reacts with water and the pH of the resultant solution is less than 7.

- (i) Write the equation to show the reaction between the salt formed and water.

.....[1]

- (ii) With reference to your equation in (b)(i), write an expression for the acid dissociation constant of the salt.

[1]

- (iii) Hence, determine the pH of the resultant solution.


[2]

[Total: 8]



- 3 Methanal is a colorless, strong-smelling gas used in making building materials and many household products.

The Strecker synthesis is a route to preparing amino acids. Glycine, 2-aminoethanoic acid, can be prepared from methanal in this way. This is shown in the four steps reaction scheme below.



- (a) (i) Suggest the role of ammonia in this synthesis.

.....[1]


- (ii) Compound **E** has the molecular formula  $\text{CH}_5\text{NO}$ . Suggest a structure for compound **E**.

[1]

- (iii) State the type of reaction for reaction 4.

.....[1]

- (b) The amino acid shown below is isoleucine, 2-amino-3-methylpentanoic acid.



2-amino-3-methylpentanoic acid


Molecule **F** can be used as the starting material to prepare this amino acid using a Strecker synthesis.

Draw the skeletal structure of **F**.

[1]

- (c) An amide bond is formed when two amino acids react together.

Phenobarbital, which is a medication used to treat epilepsy, also has an amide bond in it.




phenobarbital

Predict all the products formed when phenobarbital undergoes acidic hydrolysis.

[2]

[Total: 6]

- 4 (a) Formation of 1,2-halo alcohols, also known as halohydrins, occurs via the addition reaction between an alkene and a halogen in the presence of water.



In a series of experiments, the reaction between propene and aqueous bromine was carried out with different concentrations of the two reagents, and the following relative initial rates were obtained.

Experiment	$[\text{CH}_3\text{CH}=\text{CH}_2]$ / mol dm <sup>-3</sup>	$[\text{Br}_2]$ / mol dm <sup>-3</sup>	Initial rate / mol dm <sup>-3</sup> s <sup>-1</sup>
1	0.020	0.020	$1.00 \times 10^{-3}$
2	0.030	0.020	$1.50 \times 10^{-3}$
3	0.040	0.030	$3.00 \times 10^{-3}$

- (i) Use these data to deduce the order of reaction with respect to each of the two reagents, showing how you arrive at your answers.

[2]

- (ii) Hence write a rate equation for the reaction.

.....[1]

(iii) Calculate the rate constant for the reaction, giving its units.

[2]

(iv) State and explain how the rate of reaction may change if chlorine is used instead of bromine in the reaction with propene.

.....[1]

(b) (i) The mechanism of the addition reaction between propene and aqueous Br<sub>2</sub> involves three steps.

- There is an initial attack by the  $\pi$  electron pair of the alkene on Br<sub>2</sub> to yield a carbocation intermediate.
- This is followed by the nucleophilic attack of the lone pair of electrons on oxygen in water on the carbocation intermediate.
- The third step involves the loss of H<sup>+</sup> ion which then yields the neutral bromohydrin.

Using the information given above, describe a mechanism for this reaction.


[3]


(ii) Based on your mechanism drawn, explain whether it is consistent with the rate equation proposed.

.....  
 .....[1]

- (c) Grignard reagents are organo-magnesium halides, commonly used in synthesis to prepare a variety of organic compounds.

The carbon-magnesium bonds in Grignard reagents are highly polar and this makes it extremely useful in organic synthesis as it is able to react with other polar organic molecules to form carbon-carbon bonds. An example of the use of a Grignard reagent is the two-step reaction of  $\text{CH}_3\text{CH}_2\text{MgBr}$  with butanone,  $\text{CH}_3\text{CH}_2\text{COCH}_3$ , to form 3-methylpentan-3-ol.



- (i) Suggest the type of reaction that has taken place in step 1.  
 .....[1]
- (ii) Suggest the structural formula of the final organic product formed when  is reacted with propanone,  $\text{CH}_3\text{COCH}_3$ , in a similar two-step process.

[1]

- (iii) Suggest a suitable Grignard reagent and another organic compound to be used if propan-2-ol is to be prepared using a similar two-step process.

[2]

- (iv) The Grignard reagent  $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$  can be readily converted into a carboxylic acid by using carbon dioxide.

Suggest the structural formula for the organic product formed.

[1]

[Total: 15]

- 5 (a) Cobalt forms many coloured complexes. Cobalt(III) chloride combines with ammonia to form a pink coloured compound **A**,  $\text{CoCl}_3 \cdot \text{H}_2\text{O} \cdot 5\text{NH}_3$  ( $M_r = 268.4$ ) in which the coordination number of cobalt is 6.

1.00 g of **A** is dissolved in 25 cm<sup>3</sup> of water and the solution is titrated with 0.500 mol dm<sup>-3</sup> silver nitrate solution. It is found that 22.40 cm<sup>3</sup> of silver nitrate is required for complete reaction.

- (i) Calculate the number of moles of free chloride ions per mole of **A**.

[2]

- (ii) Draw the structure of the complex ion in **A**.

[1]

- (iii) When the pink compound **A** is heated, water vapour and ammonia were evolved to give a purple solid **B**.  
**A** and **B** have the same coordination number.

Suggest the formula of the complex in the purple solid **B**.


.....[1]

- (iv) Account for the difference in the colour of **A** and **B**.

.....  
 .....  
 .....

.....[2]

- (b) Aqueous hydrogen peroxide is fairly stable, but when a mixture of a cobalt(II) salt and tartaric acid is added to aqueous hydrogen peroxide, the initially pink solution slowly turns into a green cobalt(III) species.



After a while, oxygen gas is vigorously evolved and the solution turns back to pink again.

- (i) State the role of the cobalt(II) salt in this reaction and support your answer by referring to the observations.

Role of the cobalt(II) salt: .....[1]

Observation with explanation:

.....  
 .....  
 .....  
 .....[1]

- (ii) Tartaric acid acts as a complexing agent in this reaction to stabilise the  $\text{Co}^{3+}$  cation.

With the aid of relevant data from the *Data Booklet*, show that  $\text{Co}^{3+}$  is not stable in aqueous solution.

.....  
 .....  
 .....  
 .....[2]



(c) A student wanted to measure the standard cell potential,  $E^\ominus_{\text{cell}}$ , between the  $\text{Co}^{2+}/\text{Co}$  half-cell and the  $\text{Fe}^{3+}/\text{Fe}^{2+}$  half-cell. She set up and connected the two half cells and obtained a reading.

(i) Calculate the value  $E^\ominus_{\text{cell}}$  that will be obtained by the student.

[1]

(ii) State and explain what happens to the standard cell potential,  $E^\ominus_{\text{cell}}$  when ammonia is added to the  $\text{Co}^{2+}/\text{Co}$  half-cell.

.....

.....


.....

.....

.....[2]

[Total: 13]

6 A student plotted the sketch graph below to show how the entropy of a sample of H<sub>2</sub>O varies with temperature.



(a) Identify the state of H<sub>2</sub>O in Phase 2 and 3 respectively. Suggest a value of T<sub>2</sub>.

In phase 2:..... In phase 3: ..... value of T<sub>2</sub>: ..... [1]

(b) Suggest why entropy of H<sub>2</sub>O is zero at 0 K.

.....[1]

(c) Explain why the entropy change, ΔS, at temperature T<sub>2</sub> is much larger than that at temperature T<sub>1</sub>.

.....  
 .....  
 .....  
 .....[2]

(d) It requires 3.49 kJ of heat energy to convert 1.53 g of H<sub>2</sub>O from the state in phase 2 to phase 3 at temperature T<sub>2</sub> and 100kPa.

Use these data and your value of T<sub>2</sub> in part (a) to calculate the value of ΔS, including units, for the conversion of one mole of H<sub>2</sub>O from the state in phase 2 to phase 3 at temperature T<sub>2</sub>.

[3]

- (e) The student wants to find out if dissolving a salt, silver fluoride in water is always a spontaneous process. He must first find the enthalpy change of solution of silver fluoride in water.

Some enthalpy changes for silver fluoride are shown in the table.

	$\Delta H / \text{kJ mol}^{-1}$
Lattice energy of silver fluoride	-950
Enthalpy change of hydration for silver ions	-464
Enthalpy change of hydration for fluoride ions	-506

- (i) Use the data provided to calculate a value for the enthalpy change of solution of silver fluoride in water.

[2]

- (ii) If entropy change for dissolving silver fluoride in water has a positive value, explain why dissolving of silver fluoride in water is always a spontaneous process.

.....  
 .....  
 .....  
 .....[2]


- (iii) Explain why the enthalpy change of hydration of the fluoride ions is more negative than the enthalpy change of hydration of the chloride ions.

.....  
 .....  
 .....[1]

[Total: 12]

7 The uses of carboxylic acids are so extensive that they can be divided into several industries, such as pharmaceuticals or food among others.

(a) Ethanol and ethanoic acid react reversibly to form ethyl ethanoate and water according to the equation:



A student mixed 0.0800 mol of ethanoic acid and 0.120 mol of ethanol in a conical flask and the flask was sealed with a bung and allowed to reach equilibrium at 20 °C.

The equilibrium mixture is placed in a graduated flask and the volume made up to 250 cm<sup>3</sup> with distilled water. A 10.0 cm<sup>3</sup> sample of this equilibrium mixture was placed in a conical flask with a few drops of phenolphthalein and titrated with 0.100 mol dm<sup>-3</sup> of sodium hydroxide from a burette. The indicator turned pink when 6.40 cm<sup>3</sup> of NaOH had been added.

(i) Calculate the amount of CH<sub>3</sub>COOH in the 250 cm<sup>3</sup> equilibrium mixture.

[1]

(ii) Hence, calculate the value for  $K_c$  for the reaction of ethanoic acid and ethanol at 20 °C.

[2]

- (b) The following table compares the  $pK_a$  values of two dicarboxylic acids with that of ethanoic acid.

Acid	Formula	$pK_1$	$pK_2$
ethanoic	$\text{CH}_3\text{COOH}$	4.8	–
malonic	$\text{HOOCCH}_2\text{COOH}$	2.8	5.7
succinic	$\text{HOOC}(\text{CH}_2)_2\text{COOH}$	4.2	5.6

- (i) Suggest a reason why the  $pK_1$  value of malonic acid is so much less than the  $pK_1$  of ethanoic and succinic acid.

.....  
 .....  
 .....  
 .....[2]

- (ii) Suggest a reason why the  $pK_2$  value of malonic and succinic acid is higher than its respective  $pK_1$  value.

.....  
 .....[1]

- (c) Malonic acid can undergo dehydration with  $\text{P}_4\text{O}_{10}$  to give a foul-smelling gas, **A**. At 30.5 kPa, 0.1057 g of **A** occupies 200  $\text{cm}^3$  at a temperature of 200°C. Determine the relative molecular mass of **A**.

[2]

- (d) At high temperature, succinic acid can also undergo dehydration to produce a neutral compound **B**,  $C_4H_4O_3$  which does not react with sodium metal or Brady's reagent.


Compound **B** reacts with ammonia to give a compound **C**,  $C_4H_7NO_3$ , which reacts with cold  $NaOH(aq)$ , but not with cold  $HCl(aq)$ .

Suggest structures for **B** and **C** and explain the observations.

[4]

[Total: 12]





INNOVA JUNIOR COLLEGE  
 JC2 PRELIMINARY EXAMINATION  
 in preparation for General Certificate of Education Advanced Level  
**Higher 2**

CANDIDATE  
 NAME

CLASS

INDEX NUMBER

## CHEMISTRY

**9729/02**

Paper 2 Structured Questions

**27 August 2018**

Candidates answer on the question paper.

**2 hours**

Additional Materials: *Data Booklet*

### READ THESE INSTRUCTIONS FIRST

Write your index number, name and civics group on all the work you hand in.  
 Write in dark blue or black pen.

You may use pencil for any diagrams, graphs or rough working.  
 Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions in the space provided.

You are advised to show all working in calculations.  
 You are reminded of the need for good English and clear presentation in your answers.  
 You are reminded of the need for good handwriting.  
 Your final answers should be in 3 significant figures.

You may use a calculator.

The number of marks is given in brackets [ ]  
 at the end of each question or part question.



At the end of the examination, fasten all your work  
 securely together.

For Examiner's Use	
1	9
2	8
3	6
4	15
5	13
6	12
7	12
Significant Figures and Units	
Handwriting and Presentation	
<b>Total</b>	<b>75</b>

This document consists of **19** printed pages and **1** blank page.



1	(a)	<p>The following diagram shows the second ionisation energy of some elements.</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <caption>Second Ionisation Energy Data</caption> <thead> <tr> <th>Element</th> <th>Second Ionisation Energy / kJ mol<sup>-1</sup></th> </tr> </thead> <tbody> <tr><td>O</td><td>~3400</td></tr> <tr><td>S</td><td>~2300</td></tr> <tr><td>Cr</td><td>~1600</td></tr> <tr><td>Mn</td><td>~1500</td></tr> <tr><td>As</td><td>~1800</td></tr> <tr><td>Se</td><td>~2050</td></tr> <tr><td>Br</td><td>~2000</td></tr> <tr><td>Kr</td><td>~2400</td></tr> <tr><td>Te</td><td>~1800</td></tr> </tbody> </table>	Element	Second Ionisation Energy / kJ mol <sup>-1</sup>	O	~3400	S	~2300	Cr	~1600	Mn	~1500	As	~1800	Se	~2050	Br	~2000	Kr	~2400	Te	~1800
Element	Second Ionisation Energy / kJ mol <sup>-1</sup>																					
O	~3400																					
S	~2300																					
Cr	~1600																					
Mn	~1500																					
As	~1800																					
Se	~2050																					
Br	~2000																					
Kr	~2400																					
Te	~1800																					
		<p>(i) Explain why the second ionisation energies of elements O, S, Se and Te show a decreasing trend. [1]</p> <p>The <u>number of principal quantum shells increases</u>/ The valence electrons are <u>further away</u> [✓]  <u>It is less strongly attracted to the nucleus</u> OR <u>Less energy is required to remove the valence electrons</u> [✓], hence second ionisation energy decreases.</p>																				
		<p>(ii) Explain why the second ionisation energy of Br is lower than that of Se. [1]</p> <p>Se<sup>+</sup>: [Ar]4s<sup>2</sup>4p<sup>3</sup>                      Br<sup>+</sup>: [Ar]4s<sup>2</sup>4p<sup>4</sup></p> <p>The <u>paired 4p/np electrons</u> [✓] / 2 electrons in the same 4p orbital in Br<sup>+</sup> experience <u>interelectronic repulsion</u> [✓], less energy is required to remove an electron from this orbital.</p>																				
		<p>(iii) Write the full ground state electronic configuration for Cr. [1]</p> <p>1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>5</sup> 4s<sup>1</sup> [1]</p>																				
		<p>(iv) Draw the d<sub>xy</sub> orbital in the axes provided below. [1]</p>																				




[1]

(b) Oxygen–oxygen bond lengths in some molecules are given below:

Molecule	Bond Length
Oxygen, $O_2$	0.121 nm
Hydrogen peroxide, $H_2O_2$	0.149 nm
Ozone, $O_3$	0.128 nm




(i) Draw the structure of the molecule ozone,  $O_3$ .




[1]



[1]





		(ii)	With reference to the data given above, comment and explain the oxygen–oxygen bond length in ozone as compared to oxygen and hydrogen peroxide. [2]
			Oxygen–oxygen bond length in ozone is longer than O=O in oxygen but shorter than O-O in hydrogen peroxide. [1]  P orbital of oxygen overlap with $\pi$ orbital of adjacent oxygen-oxygen double bond. OR Lone pair on oxygen atom can delocalised into the $\pi$ electron cloud of adjacent oxygen- oxygen double bond [1]
		(iii)	Explain clearly why hydrogen peroxide is a liquid while oxygen is a gas at room conditions in terms of structure and bonding. [2]
			Both hydrogen peroxide and oxygen have simple molecular structure. [✓]  Oxygen have instantaneous dipole induced dipole forces of attraction between molecules [✓] while hydrogen peroxide have hydrogen bonding between the molecules. [✓] More energy is needed to overcome the stronger hydrogen bond [✓], hence it has stronger intermolecular forces of attraction and exist as a liquid.
			[Total: 9]



2	Compounds containing nitrogen are important to life and have applications in science and medicine.  Three nitrogeneous bases have the following structures.		
	 <p>N,4-dimethylbenzenamine</p>	 <p>4-methylphenylamine</p>	 <p>benzylamine</p>
(a)	(i)	Arrange the three bases above in increasing order of $pK_b$ . [1]	
		Lowest $pK_b$ $\longrightarrow$ Highest $pK_b$	


			 <p>benzylamine      <i>N</i>,4-dimethyl benzenamine      4-methylphenylamine [1]</p>
	(ii)	Explain in terms of their molecular structures why benzylamine and 4-methylphenylamine have different $pK_b$ values.	[2]
		<p>In benzylamine, the presence of <u>electron donating</u>  <u>increases the electron density on N</u> ✓, making the <u>lone pair of electron on N more available</u> ✓ to accept a proton. Hence, benzylamine is the most basic.</p> <p>In 4-methylphenylamine, the <u>lone pair of electron on N delocalised into the <math>\pi</math> orbital of the benzene ring, decreasing the electron density on N</u> ✓, making the <u>lone pair of electron on N less available</u> ✓ to accept a proton. Hence, 4-methylphenylamine is the least basic.</p>	
	(iii)	Outline how <i>N</i> ,4-dimethylbenzenamine may be produced from 4-methylphenylamine.	[1]
		Heat 4-methylphenylamine with (ethanolic) $\text{CH}_3\text{Cl}$ (in a sealed tube).	[1]
	(b)	The $pK_b$ of benzylamine is 4.66.	
		25.0 $\text{cm}^3$ of 0.025 $\text{mol dm}^{-3}$ benzylamine was completely neutralised by dilute hydrochloric acid of the same concentration. The salt formed reacts with water and the pH of the resultant solution is less than 7.	
	(i)	Write the equation to show the reaction between the salt formed and water.	[1]
		 <p>[1]</p>	

		(ii)	With reference to your equation in (b)(i), write an expression for the acid dissociation constant of the salt. <span style="float: right;">[1]</span>
			$K_a = \frac{\left[ \text{C}_6\text{H}_5\text{CH}_2\text{NH}_2 \right] \left[ \text{H}_3\text{O}^+ \right]}{\left[ \text{C}_6\text{H}_5\text{CH}_2\text{NH}_3^+ \right]}$ <span style="float: right;">[1]</span>
		(iii)	Hence, determine the pH of the resultant solution. <span style="float: right;">[2]</span>
			$[\text{salt}] = (6.25 \times 10^{-4}) \div (50/1000)$ $= 0.0125 \text{ mol dm}^{-3} \quad \checkmark$ $K_a = K_w / K_b$ $= \frac{10^{-14}}{10^{-4.66}}$ $= 4.571 \times 10^{-10} \quad \checkmark$ $[\text{H}^+] = \sqrt{K_a \times [\text{salt}]}$ $= \sqrt{4.571 \times 10^{-10} \times 0.0125}$ $= 2.390 \times 10^{-6} \quad \checkmark$ $\text{pH} = 5.62 \quad \checkmark$
			[Total: 8]

<b>3</b>	<p>Methanal is a colorless, strong-smelling gas used in making building materials and many household products.</p> <p>The Strecker synthesis is a route to preparing amino acids. Glycine, 2-aminoethanoic acid, can be prepared from methanal in this way. This is shown in the four steps reaction scheme below.</p>
----------	--

	 <p style="text-align: center;">2-aminoethanoic acid</p>	
(a)	(i)	<p>What is the role of ammonia, in the context of this synthesis? [1]</p> <p>Nucleophile [1] Allow Nucleophilic or Lewis base or Lone-pair donor</p>
	(ii)	<p>Compound <b>E</b> has the molecular formula <math>\text{CH}_5\text{NO}</math>. Suggest a structure for compound <b>E</b>. [1]</p> <p> [1]</p>
	(iii)	<p>State the type of reaction for reaction 4. [1]</p> <p>Acidic hydrolysis [1]</p>
(b)	<p>The amino acid shown below is isoleucine, 2-amino-3-methylpentanoic acid.</p> <p style="text-align: center;">  </p> <p>Molecule <b>F</b> can be used as the starting material to prepare this amino acid using a Strecker synthesis. Draw the skeletal structure of <b>F</b>. [1]</p> <p> [1]</p>	

	(c)	<p>An amide bond is formed when two amino acids react together.</p> <p>Phenobarbital, which is a medication used to treat epilepsy, also has an amide bond in it.</p>  <p style="text-align: center;">phenobarbital</p> <p>Predict all the products formed when phenobarbital undergoes acidic hydrolysis.</p>	
		 <p style="text-align: center;">[1] , CO<sub>2</sub> [✓], NH<sub>4</sub><sup>+</sup> [✓]</p>	
			[2]
			[Total: 6]

4	(a)	<p>Formation of 1,2-halo alcohols, also known as halohydrins, occurs via the addition reaction between an alkene and a halogen in the presence of water.</p>  <p style="text-align: center;">halohydrin</p> <p>In a series of experiments, the reaction between propene and aqueous bromine was carried out with different concentrations of the two reagents, and the following relative initial rates were obtained.</p>
---	-----	--

Experiment	[CH <sub>3</sub> CH=CH <sub>2</sub> ] / mol dm <sup>-3</sup>	[Br <sub>2</sub> ] / mol dm <sup>-3</sup>	initial rate / mol dm <sup>-3</sup> s <sup>-1</sup>
1	0.020	0.020	1.00 x 10 <sup>-3</sup>
2	0.030	0.020	1.50 x 10 <sup>-3</sup>
3	0.040	0.030	3.00 x 10 <sup>-3</sup>


(i)	Use these data to deduce the order of reaction with respect to each of the two reagents, showing how you arrive at your answers. [2]		
	Let the rate equation be: Rate = k[CH <sub>3</sub> CH=CH <sub>2</sub> ] <sup>x</sup> [Br <sub>2</sub> ] <sup>y</sup>		
	<table border="1"> <tr> <td> <p>Comparing experiments 1 &amp; 2,</p> <math display="block">\frac{\text{Rate}_1}{\text{Rate}_2} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}</math> <math display="block">\frac{1.00 \times 10^{-3}}{1.50 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.030]^x [0.020]^y}</math> <p>x = 1 [1m]</p> </td> <td> <p>Comparing experiments 1 &amp; 3,</p> <math display="block">\frac{\text{Rate}_1}{\text{Rate}_3} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}</math> <math display="block">\frac{1.00 \times 10^{-3}}{3.00 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.040]^x [0.030]^y}</math> <p>y = 1 [1m]</p> </td> </tr> </table>	<p>Comparing experiments 1 &amp; 2,</p> $\frac{\text{Rate}_1}{\text{Rate}_2} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}$ $\frac{1.00 \times 10^{-3}}{1.50 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.030]^x [0.020]^y}$ <p>x = 1 [1m]</p>	<p>Comparing experiments 1 &amp; 3,</p> $\frac{\text{Rate}_1}{\text{Rate}_3} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}$ $\frac{1.00 \times 10^{-3}}{3.00 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.040]^x [0.030]^y}$ <p>y = 1 [1m]</p>
<p>Comparing experiments 1 &amp; 2,</p> $\frac{\text{Rate}_1}{\text{Rate}_2} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}$ $\frac{1.00 \times 10^{-3}}{1.50 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.030]^x [0.020]^y}$ <p>x = 1 [1m]</p>	<p>Comparing experiments 1 &amp; 3,</p> $\frac{\text{Rate}_1}{\text{Rate}_3} = \frac{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}{k [\text{CH}_3\text{CH} = \text{CH}_2]^x [\text{Br}_2]^y}$ $\frac{1.00 \times 10^{-3}}{3.00 \times 10^{-3}} = \frac{k [0.020]^x [0.020]^y}{k [0.040]^x [0.030]^y}$ <p>y = 1 [1m]</p>		
(ii)	Hence write a rate equation for the reaction. [1]		
	Rate = k[CH <sub>3</sub> CH=CH <sub>2</sub> ][Br <sub>2</sub> ]		
(iii)	Calculate the rate constant for the reaction, giving its units. [2]		
	<p>Using experiment 1,</p> $k = \frac{\text{Rate}}{[\text{CH}_3\text{CH} = \text{CH}_2][\text{Br}_2]}$ $k = \frac{1.00 \times 10^{-3}}{[0.020][0.020]} = 2.50 [1m] \text{ mol}^{-1}\text{dm}^3\text{s}^{-1} [1m]$		
(iv)	State and explain how the rate of reaction may change if chlorine is used instead of bromine in the reaction with propene. [1]		
	Rate will decrease [✓] because C-Cl bond is stronger than Br-Br [✓].		
(b) (i)	<p>The mechanism of the addition reaction between propene and aqueous Br<sub>2</sub> involves three steps.</p> <ul style="list-style-type: none"> <li>There is an initial attack by the π electron pair of the alkene on Br<sub>2</sub> to yield a carbocation intermediate.</li> </ul>		




- This is followed by the nucleophilic attack of the lone pair of electrons on oxygen in water on the carbocation intermediate.
- The third step involves the loss of  $H^+$  ion which then yields the neutral bromohydrin.

Using the information given above, describe a mechanism for this reaction.


[3]



Dipoles [ $\checkmark$ ], two curly arrows [ $\checkmark$ ], correct carbocation [ $\checkmark$ ], label slow step [ $\checkmark$ ],  $Br^-$  [ $\checkmark$ ]



Lone pair + curly arrow [ $\checkmark$ ], intermediate [ $\checkmark$ ], label fast step [ $\checkmark$ ]



Curvy arrow [ $\checkmark$ ], final organic product [ $\checkmark$ ],  $H^+$  [ $\checkmark$ ]

11 [ $\checkmark$ ] – 3m

6 to 10 [ $\checkmark$ ] – 2m

3 to 5 [ $\checkmark$ ] – 1m

-

- (ii) Based on your mechanism drawn, explain whether it is consistent with the rate equation proposed. [1]


It is **consistent** as there is **1 propene molecule and 1  $Br_2$  molecule in the rate determining step**, which is consistent with the rate equation.

Or

To allow ecf if rate equation is wrong

(c) Grignard reagents are organo-magnesium halides, commonly used in synthesis to prepare a variety of organic compounds.


The carbon-magnesium bonds in Grignard reagents are highly polar and this makes it extremely useful in organic synthesis as it is able to react with other polar organic molecules to form carbon-carbon bonds. An example of the use of a Grignard reagent is the two-step reaction of  $\text{CH}_3\text{CH}_2\text{MgBr}$  with butanone,  $\text{CH}_3\text{CH}_2\text{COCH}_3$ , to form 3-methylpentan-3-ol.




(i) Suggest the type of reaction that has taken place in step 1. [1]

Nucleophilic addition [1]


(ii) Suggest the structural formula of the final organic product formed when




is reacted with propanone,  $\text{CH}_3\text{COCH}_3$ , in a similar two-step process. [1]



		(iii)	Suggest a suitable Grignard reagent and another organic compound to be used if propan-2-ol is to be prepared using a similar two-step process. [2]
			$\text{CH}_3\text{MgBr}$ [1] $\text{CH}_3\text{CHO}$ [1]
		(iv)	The Grignard reagent $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$ can be readily converted into a carboxylic acid by using carbon dioxide. Suggest the structural formula for the organic product formed. [1]
			$\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$
			[Total: 15]

5	(a)	<p>Cobalt forms many coloured complexes. Cobalt(III) chloride combines with ammonia to form a pink coloured compound <b>A</b>, <math>\text{CoCl}_3 \cdot \text{H}_2\text{O} \cdot 5\text{NH}_3</math> (<math>M_r = 268.4</math>) in which the coordination number of cobalt is 6.</p> <p>1.00 g of <b>A</b> is dissolved in <math>25 \text{ cm}^3</math> of water and the solution is titrated with <math>0.500 \text{ mol dm}^{-3}</math> silver nitrate solution. It is found that <math>22.40 \text{ cm}^3</math> of silver nitrate is required for complete reaction.</p>
	(i)	<p>Calculate the number of moles of free chloride ions per mole of <b>A</b>.</p> <p>no. of moles of <math>\text{AgNO}_3 = 0.500 \times (22.40/1000) = 1.12 \times 10^{-2} \text{ mol}</math> [1]  no. of moles of <math>\text{Cl}^- = 1.12 \times 10^{-2} \text{ mol}</math>  no. of moles of <b>A</b> = <math>1.00/268.4 = 3.73 \times 10^{-3} \text{ mol}</math>  no. of moles of free <math>\text{Cl}^-</math> ions per mole of <b>A</b> = <math>(1.12 \times 10^{-2}) / (3.73 \times 10^{-3}) = 3</math> [1]</p>
	(ii)	<p>Draw the structure of the complex ion in <b>A</b>.</p> 
	(iii)	<p>When the pink compound <b>A</b> is heated, water vapour and ammonia were evolved to give a purple solid <b>B</b>.  <b>A</b> and <b>B</b> have the same coordination number.  Suggest the formula of the complex in the purple solid <b>B</b>.</p>
		$[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$ or $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$ [1]

			[1]
	(iv)	Account for the difference in the colour of <b>A</b> and <b>B</b> .	
		The <b>different ligands</b> in <b>A</b> and <b>B</b> cause the <b>energy gap between the split d-orbitals to be different</b> . [1] The <b>wavelength of light absorbed is different</b> [1] for <b>A</b> and <b>B</b> and thus the colour observed are different.	[2]
	(b)	<p>Aqueous hydrogen peroxide is fairly stable, but when a mixture of a cobalt(II) salt and tartaric acid is added to aqueous hydrogen peroxide, the initially pink solution slowly turns into a green cobalt(III) species.</p> <div style="text-align: center;">  <p>tartaric acid</p> </div> <p>After a while, oxygen gas is vigorously evolved and the solution turns back to pink again.</p>	
	(i)	State the role of the cobalt(II) salt in this reaction and support your answer by referring to the observations.	
		<p>Role of the cobalt(II) salt: <b>catalyst</b> [1]</p> <p>Observation with explanation:</p> <ul style="list-style-type: none"> <li>The <b>oxygen gas is vigorously evolved</b> showing that the <b>reaction speeds up</b> [1] OR</li> <li>The solution turns from <b>pink to green to pink (WTTE)</b> showing that <b>Co<sup>2+</sup> is regenerated</b> [1]</li> </ul>	[2]
	(ii)	<p>Tartaric acid acts as a complexing agent in this reaction to stabilise the Co<sup>3+</sup> cation.</p> <p>With the aid of relevant data from the <i>Data Booklet</i>, show that Co<sup>3+</sup> is not stable in aqueous solution.</p>	
		$\text{Co}^{3+} + e \rightleftharpoons \text{Co}^{2+} \quad E_{\text{red}} = +1.89\text{V}$ $\text{O}_2 + 4\text{H}^+ + 4e \rightleftharpoons 2\text{H}_2\text{O} \quad E_{\text{red}} = +1.23\text{V}$ $E^\circ_{\text{cell}} = +1.89 - (+1.23) = +0.66\text{V} \quad [1]$ <p>Co<sup>3+</sup> oxidises water readily to form Co<sup>2+</sup> / Co<sup>3+</sup> is readily reduced by water to form Co<sup>2+</sup>. [1]</p>	[2]
	(c)	A student wanted to measure the standard cell potential, $E^\circ_{\text{cell}}$ , between the Co <sup>2+</sup> /Co half-cell and the Fe <sup>3+</sup> /Fe <sup>2+</sup> half-cell. She set up and connected the two half cells and obtained a reading.	
	(i)	Calculate the value $E^\circ_{\text{cell}}$ that will be obtained by the student.	
	[R]	$\text{Fe}^{3+} + e \rightleftharpoons \text{Fe}^{2+} \quad E_{\text{red}} = +0.77\text{V}$	
	[O]	$\text{Co}^{2+} + 2e \rightleftharpoons \text{Co} \quad E_{\text{red}} = -0.28\text{V}$	

		$E^\ominus_{\text{cell}} = +0.77 - (-0.28) = +1.05 \text{ V}$ [1]	[1]
	(ii)	State and explain what happens to the standard cell potential, $E^\ominus_{\text{cell}}$ when ammonia is added to the $\text{Co}^{2+}/\text{Co}$ half-cell.	
		<p>When <math>\text{NH}_3</math> is added, the following equilibrium is set up in the <math>\text{Co}^{2+}/\text{Co}</math> half-cell:</p> <p>Either:</p> <p><math>[\text{Co}(\text{NH}_3)_6]^{2+} + 2e \rightleftharpoons \text{Co} + 6\text{NH}_3</math> <math>E_{\text{red}} = -0.43\text{V}</math> [1]  (Give B.O.D. if the value <math>-0.43\text{V}</math> is not quoted in the answer)</p> <p><b>OR <math>[\text{Co}^{2+}]</math> decreases</b> as <math>\text{Co}(\text{OH})_2(\text{s})</math> is formed. [1]</p> <p>the <math>E_{\text{ox}}</math> of <math>\text{Co}^{2+}/\text{Co}</math> <b>will become more negative/ position of the equilibrium</b>  <math>\text{Co}^{2+} + 2e \rightleftharpoons \text{Co}</math> <b>will shift left</b></p> <p>Hence, the <math>E^\ominus_{\text{cell}}</math> becomes <b>more positive</b>. [1]</p>	[2]
			[Total: 13]

6	A student plotted the sketch graph below to show how the entropy of a sample of $\text{H}_2\text{O}$ varies with temperature.
(a)	Identify the state of $\text{H}_2\text{O}$ in Phase 2 and 3 respectively. Suggest a value of $T_2$ .
	In phase 2:..... In phase 3: ..... value of $T_2$ : ..... [1]
	In phase 2: liquid In phase 3: gas $T_2$ : 373K
(b)	Suggest why entropy of $\text{H}_2\text{O}$ is zero at 0 K.
	.....[1]
	It is at maximum state of order or no disorder or perfectly ordered or pure crystalline structure.

(c)	Explain why the entropy change, $\Delta S$ , at temperature $T_2$ is much larger than that at temperature $T_1$ .								
	<p>.....</p> <p>.....</p> <p>.....</p> <p>.....</p> <p>.....</p> <p style="text-align: right;">[2]</p>								
	<p>At <math>T_2</math>, the increase from phase 2 to 3 is much larger as the <u>gas molecules</u> in phase 3 have <u>more ways in which the particles can be arranged</u> than particles in liquid. [1]</p> <p>At <math>T_1</math>, the particles in phase 2 (liquid) are still orderly arranged compared to the ordered arrangement of particles in phase 1. [1] Thus, the increase in disorder is smaller from phase 1 to 2. (2<sup>nd</sup> marking point: mention of smaller increase from phase 1 to 2 as it is from an ordered arrangement to slightly disordered arrangement in liq.)</p>								
(d)	<p>It requires 3.49 kJ of heat energy to convert 1.53 g of <math>H_2O</math> from the state in phase 2 to phase 3 at temperature <math>T_2</math> and 100kPa.</p> <p>Use these data and your value of <math>T_2</math> in part (a) to calculate the value of <math>\Delta S</math>, including units, for the conversion of one mole of <math>H_2O</math> from the state in phase 2 to phase 3 at temperature <math>T_2</math>.</p>								
	<p style="text-align: right;">[3]</p> <p>Amt of water = <math>\frac{1.53}{18} = 0.0850</math></p> <p>enthalpy change reaction per mole of water = <math>\frac{3.49}{0.0850} = +41.1 \text{ kJ mol}^{-1}</math> [1] award if + sign is omitted, but penalise negative sign is included</p> <p><math>\Delta G = \Delta H - T\Delta S = 0</math> <math>+41.1 - (373)\Delta S = 0</math>      equate <math>\Delta G = 0</math>, substitute <math>\Delta H</math>, <math>T_2</math> into equation [1] ecf</p> <p><math>\Delta S = 0.110 \text{ kJ K}^{-1} \text{ mol}^{-1}</math>    correct computation (allow ecf) and units [1] ecf</p>								
(e)	<p>The student wants to find out if dissolving a salt, silver fluoride in water is always a spontaneous process. He must first find the enthalpy change of solution of silver fluoride in water.</p> <p>Some enthalpy changes for silver fluoride are shown in the table.</p> <table border="1" data-bbox="303 1713 1332 1926"> <thead> <tr> <th></th> <th><math>\Delta H / \text{kJ mol}^{-1}</math></th> </tr> </thead> <tbody> <tr> <td>Lattice energy of silver fluoride</td> <td>-950</td> </tr> <tr> <td>enthalpy change of hydration for silver ions</td> <td>-464</td> </tr> <tr> <td>enthalpy change of hydration for fluoride ions</td> <td>-506</td> </tr> </tbody> </table>		$\Delta H / \text{kJ mol}^{-1}$	Lattice energy of silver fluoride	-950	enthalpy change of hydration for silver ions	-464	enthalpy change of hydration for fluoride ions	-506
	$\Delta H / \text{kJ mol}^{-1}$								
Lattice energy of silver fluoride	-950								
enthalpy change of hydration for silver ions	-464								
enthalpy change of hydration for fluoride ions	-506								
(i)	Use the data provided to calculate a value for the enthalpy change of solution of silver fluoride in water.								

			[2]
		$\Delta H_{\text{solution}} = \Delta H_{\text{hyd}} \text{Ag}^+ + \Delta H_{\text{hyd}} \text{F}^- - LE$ $= (-464) + (-506) - (-950) \quad [1] \text{ working}$ $= -20 \text{ kJ mol}^{-1} \quad [1] \text{ answer}$	
	(ii)	If entropy change for dissolving silver fluoride in water has a positive value, explain why dissolving of silver fluoride in water is always a spontaneous process.	
			[2]
		$\Delta G = \Delta H - T\Delta S$ <p>Relate to signs of <math>\Delta S</math> and <math>\Delta H</math> [1]:          Since <math>\Delta S</math> is positive, the term <math>(-T\Delta S)</math> is always negative.          Thus, since <math>\Delta H</math> is negative, <u><math>\Delta G</math> is always negative at all temperature.</u> [1]</p>	
	(iii)	Explain why the enthalpy change of hydration of the fluoride ions is more negative than the enthalpy change of hydration of the chloride ions.	
			[1]
		$\Delta H_{\text{hydration}} \propto \text{charge density} [\checkmark]$ Fluoride (ions) are smaller (than chloride), with <u>higher charge density</u> $[\checkmark]$	
			[Total: 12]

<b>7</b>	The uses of carboxylic acids are so extensive that they can be divided into several industries, such as pharmaceuticals or food among others.
<b>(a)</b>	<p>Ethanol and ethanoic acid react reversibly to form ethyl ethanoate and water according to the equation:</p> $\text{CH}_3\text{COOH}(\text{l}) + \text{CH}_3\text{CH}_2\text{OH}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOCH}_2\text{CH}_3(\text{l}) + \text{H}_2\text{O}(\text{l})$ <p>A student mixed <math>8.00 \times 10^{-2}</math> mol of ethanoic acid and <math>1.20 \times 10^{-1}</math> mol of ethanol in a conical flask and the flask was sealed with a bung and allowed to reach equilibrium at <math>20^\circ\text{C}</math>.</p> <p>The equilibrium mixture is placed in a graduated flask and the volume made up to <math>250 \text{ cm}^3</math> with distilled water. A <math>10.0 \text{ cm}^3</math> sample of this equilibrium mixture was placed in a conical flask with a few drops of phenolphthalein and titrated with <math>0.100 \text{ mol dm}^{-3}</math> of sodium hydroxide from a burette. The indicator turned pink when <math>6.40 \text{ cm}^3</math> of NaOH had been added.</p>
<b>(i)</b>	<p>Calculate the amount of <math>\text{CH}_3\text{COOH}</math> in the equilibrium mixture.</p> <p>Amount of NaOH = <math>0.100 \times \frac{6.40}{1000} = 0.00064 \text{ mol}</math></p> <p>Amount of <math>\text{CH}_3\text{COOH}</math> in <math>10.0 \text{ cm}^3 = 0.00064 \text{ mol}</math></p> <p>Amount of <math>\text{CH}_3\text{COOH}</math> in <math>250 \text{ cm}^3 = \frac{250}{10} \times 0.00064 = 0.016 \text{ mol}</math> [1]</p>

[1]

- (ii) Hence, calculate the value for  $K_c$  for the reaction of ethanoic acid and ethanol at 20 °C.

	CH <sub>3</sub> CO <sub>2</sub> H (l)	C <sub>2</sub> H <sub>5</sub> OH (l)	CH <sub>3</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> (l)	H <sub>2</sub> O (l)
Initial amt / mol	0.08	0.12	0	0
Equilibrium amt / mol	<b>0.016</b>	<b>0.056</b>	<b>0.064</b>	<b>0.064</b>

Equilibrium amount [1]

$$K_c = \frac{[\text{H}_2\text{O}][\text{CH}_3\text{COOC}_2\text{H}_5]}{[\text{CH}_3\text{COOH}][\text{C}_2\text{H}_5\text{OH}]}$$

$$= \frac{\left(\frac{0.064}{V}\right)\left(\frac{0.064}{V}\right)}{\left(\frac{0.016}{V}\right)\left(\frac{0.056}{V}\right)}$$

$$= 4.57 \text{ [1]}$$

[2]

- (b) The following table compares the  $pK_a$  values of two dicarboxylic acids with that of ethanoic acid.

acid	Formula	$pK_1$	$pK_2$
ethanoic	CH <sub>3</sub> COOH	4.8	–
malonic	HOOCCH <sub>2</sub> COOH	2.8	5.7
succinic	HOOC(CH <sub>2</sub> ) <sub>2</sub> COOH	4.2	5.6

- (i) Suggest a reason why the  $pK_1$  value of malonic acid is so much less than the  $pK_1$  of ethanoic and succinic acid.

Malonic acid is **more acidic** than ethanoic or succinic acid [1] as its **monoanion is stabilised by intra hydrogen bonds between the COO<sup>-</sup> group and unionised COOH group** [1].



[2]


- (ii) Suggest a reason why the  $pK_2$  value of malonic and succinic acid is higher than its respective  $pK_1$  value.

The removal of an H<sup>+</sup> from the monoanion that already carries a negative charge would be electrostatically unfavourable. [1]

[1]



(c)	Malonic acid can undergo dehydration with $P_4O_{10}$ to give a foul-smelling gas, <b>A</b> . At 30.5 kPa, 0.1057 g of <b>A</b> occupies 200 cm <sup>3</sup> at a temperature of 200°C. Determine the relative molecular mass of <b>A</b> .
	$pV = nRT$ $30.5 \times 10^3 \times \frac{200}{10^6} = \frac{0.1057}{M_r} \times 8.31 \times (200 + 273)$ [1] $M_r = 68.1$ (1 d.p) [1]
	[2]
(d)	<p>At high temperature, succinic acid can also undergo dehydration to produce a neutral compound <b>B</b>, <math>C_4H_4O_3</math> which does not react with sodium metal or Brady's reagent.</p> <p>Compound <b>B</b> reacts with ammonia to give a compound <b>C</b>, <math>C_4H_7NO_3</math>, which reacts with cold NaOH(aq), but not with cold HCl(aq).</p> <p>Suggest structures for <b>B</b> and <b>C</b> and explain the observations.</p>
	<ul style="list-style-type: none"> <li>• <b>B</b> does not react with Na; <b>absence of –OH group</b>. [✓]</li> <li>• <b>B</b> does not react with Brady's reagent; <b>absence of carbonyl group</b>. [✓]</li> <li>• <b>C</b> does not react with cold HCl(aq); <b>absence of basic group</b> [✓], <b>amide</b> likely to be present [✓].</li> <li>• <b>C</b> undergoes <b>acid base</b> with cold NaOH(aq) [✓]; it contains an <b>acidic group</b> (carboxylic acid). [✓]</li> </ul> <p><b>Max [2] for explanations</b></p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>compound B</p> </div> <div style="text-align: center;">  <p>compound C</p> </div> </div> <p><b>[1] each</b></p>
	[4]
	[Total: 12]



INNOVA JUNIOR COLLEGE  
JC 2 PRELIMINARY EXAMINATION  
in preparation for General Certificate of Education Advanced Level  
**Higher 2**

---

**CHEMISTRY**

**9729/03**

Paper 3 Free Response

**10 September 2018**

**2 hours**

Candidates answer on separate paper.

Additional Materials: Writing Papers  
*Data Booklet*  
Cover Page

---

**READ THESE INSTRUCTIONS FIRST**

Write your name and class on all the work you hand in.  
Write in dark blue or black pen on both sides of the paper.  
You may use a soft pencil for any diagrams, graphs or rough working.  
Do not use staples, paper clips, highlighters, glue or correction fluid.

**Section A**

Answer **all** questions.

**Section B**

Answer **one** question.

You are advised to show all working in calculations.  
You are reminded of the need for good English and clear presentation in your answers.  
You are reminded of the need for good handwriting.  
Your final answers should be in 3 significant figures.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in the brackets [ ] at the end of each question or part question.

---

This document consists of **13** printed pages and **1** blank page.




**Section A**


Answer **all** the questions in this section.

- 1 This question is about the chemistry of ethene and its derivatives.
- (a) Ethene is the starting material to form ethanedioic acid.
- (i) Suggest the synthetic route for the formation of ethanedioic acid from ethene. [2]
- (ii) 0.200 mol of ethene is stored in a 20.0 dm<sup>3</sup> flask with 0.800 mol of ethane at 127 °C.
- Calculate the total pressure in the flask. Hence or otherwise, calculate the partial pressure of ethene in the flask. [2]
- (iii) The total pressure that you calculated in (a)(ii) is different from the actual pressure exerted. Suggest an explanation for the difference. [1]
- (iv) Ethene reacts with hydrogen in the presence of nickel catalyst to form ethane.
- Explain why nickel can be used as a catalyst in this reaction. [2]
- (v) Outline the mode of action of the catalyst in this reaction. [2]
- (b) Dissolving  $4.82 \times 10^{-5}$  mol calcium ethanedioate, CaC<sub>2</sub>O<sub>4</sub>, in 1 dm<sup>3</sup> of water forms a saturated solution.
- (i) Write an expression for the solubility product of calcium ethanedioate and state its units. [1]
- (ii) Calculate the solubility product of calcium ethanedioate. [1]
- (iii) 50.0 cm<sup>3</sup> of 0.100 mol dm<sup>-3</sup> CaCl<sub>2</sub> and 50.0 cm<sup>3</sup> of 0.300 mol dm<sup>-3</sup> Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub> are mixed together. Determine if CaC<sub>2</sub>O<sub>4</sub> will be precipitated out. [2]

- (c) Synthesis of the unionised form of EDTA ( $H_4Y$ ) can be carried out in the laboratory using ethene as the starting material.



- (i) Draw the structures of **A** and **B**. [2]
- (ii) Suggest appropriate reagents and conditions for step 3. [1]
- (d) Ethene can be used to form ethanal. Ethanal in turn is used to synthesize glycolic acid via the reaction shown below.



- (i) Identify the intermediate **C** and state the reagent and condition for step 3. [2]
- (ii) Explain the difference in acidity between glycolic acid and ethanoic acid. [2]
- (iii) Two molecules of glycolic acid can react with one another under suitable conditions to form a neutral compound with the loss of two water molecules.
- Suggest a possible structure for the compound formed. [1]
- (iv) Suggest a simple chemical test to distinguish between ethanoic acid and glycolic acid. [2]

[Total: 23]

- 2 Iron is the fourth most common element in the Earth's crust. It is a d-block element which is known to exhibit different characteristics from the s-block elements. Since ancient times, iron has been widely employed in a variety of applications.

- (a) (i) One well-known property of iron and its compounds is the ability to catalyse reactions. For example, aqueous iron(II) chloride can be used to catalyse the reaction between  $I^-$  and  $S_2O_8^{2-}$ , to form  $I_2$  and  $SO_4^{2-}$  ions.

Using relevant  $E^\ominus$  values from the *Data Booklet*, explain why iron(II) chloride can be used as a catalyst for this reaction.


[2]

- (ii) A sample of iron was vapourised, ionised and passed through an electric field. Analysis of deflection occurring at the electric field region revealed that a sample of  $^{32}S^{2-}$  ions would be deflected by  $+20^\circ$  towards the positive potential.

What is the angle, and direction of deflection for a sample of  $^{56}Fe^{3+}$  ions passing through the same electric field?

[2]

- (b) Ferrocene,  $Fe(C_5H_5)_2$ , is an orange solid which is known to exhibit anti-cancer activity. In this complex,  $C_5H_5^-$  is the ligand and it donates  $\pi$  electrons from the ring to the vacant 3d orbital of Fe. The structure of ferrocene is given below.



ferrocene


- (i) State the oxidation number of Fe in ferrocene. [1]
- (ii) Suggest why ferrocene is a coloured complex. [2]
- (iii) Light of a longer wavelength is lower in energy than light of a shorter wavelength. The following table shows the spectral colours and the corresponding wavelengths.

Colour	Wavelength / nm
Violet	380 – 450
Blue	450 – 495
Green	495 – 570
Yellow	570 – 590
Orange	590 – 620
Red	620 – 750

Given that aqueous  $Fe^{2+}$  ion is green in colour, suggest and explain if water causes a larger split between the two groups of 3d orbitals as compared to  $C_5H_5^-$ .

[2]

- (c) Ethylbenzene is used to synthesize (3-chlorophenyl)methanol which is used as a general solvent for inks, paints, lacquers, epoxy resin coatings and as a degreasing agent.



Starting with ethylbenzene, outline a three-step reaction scheme to obtain (3-chlorophenyl)methanol. Your answer should include clearly the reagents and conditions in each step, and the structures of all intermediates formed.

[3]

- (d) **D** is an achiral organic compound with the molecular formula  $C_2H_7NO$ . It can be formed from the reaction between a primary amide and lithium aluminium hydride. In the presence of a suitable catalyst, 1 mole of **D** reacts with 1 mole of benzoic acid to form **E**,  $C_9H_{11}NO_2$ . However, 1 mole of **D** requires 2 moles of benzoyl chloride to react completely to form **F**,  $C_{16}H_{15}NO_3$  and copious white fumes. 2 moles of **D** can also react with gaseous  $PCl_5$  to form a cyclic **G**,  $C_4H_{10}N_2$ , which contains a 6-membered ring.



Deduce the structures of **D**, **E**, **F** and **G** and explain the reactions described.


[7]

[Total: 19]


- 3 An organic radical battery (ORB) is a relatively new type of battery which uses flexible plastics, to provide electrical power. One type of hybrid ORB/Li-ion battery consists of: a cathode made from solid organic polymer containing oxoammonium ion formed from (2,2,6,6-Tetramethylpiperidin-1-yl)oxyl (TEMPO) nitroxide radicals; an anode made from graphite with Li atoms inserted between the graphite layers; and an electrolyte of  $\text{LiPF}_6$  dissolved in organic solvent.

During discharge, Li atoms give up electrons at the anode to become  $\text{Li}^+$  ions. The electrons travel round the external circuit, and are picked up by the cathode. The anions and cations in the electrolyte move to the anode and cathode respectively. This is illustrated in the following

diagram in which  and  are simplified representations of the polymer containing oxoammonium ion and TEMPO nitroxide radicals respectively.



- (a) (i) Graphite is often mixed in the polymer used for making the cathode electrode.  
Suggest a reason for this. [1]
- (ii) Calculate the oxidation number of N in the cathode **before** discharge. [1]
- (iii) The  $E_{\text{cell}}$  generated by the hybrid ORB/Li-ion battery under standard conditions is 2.17 V.  
Use relevant  $E^\circ$  value from the *Data Booklet* to calculate the electrode potential generated by the cathode half-cell. [1]
- (iv) During discharge, the following reaction occurs at the cathode.



where R represents the organic polymer cathode.

Write an equation for the overall process that occurs during discharge. [1]

- (v) Draw the dot-and-cross diagram of the  $\text{PF}_6^-$  ion and state its shape. [2]
- (vi) Suggest whether  $\text{LiPF}_6$  or  $\text{LiF}$  has a lower melting point. Explain your answer. [2]
- (b) The hybrid ORB/Li-ion battery is a secondary battery, i.e., it is rechargeable.

During charging, 1.22 g of Li is regenerated from  $\text{Li}^+$  ions at the cathode.

- (i) Calculate the amount of electrons required to form 1.22 g of Li. [1]

Besides the generation of Li, there is a competing side-reaction that occurs at the cathode.

In this side-reaction, ethylene carbonate,  $\text{C}_3\text{H}_4\text{O}_3$  undergoes reduction in the presence of  $\text{Li}^+$  ions to form ethene and lithium carbonate.


- (ii) Write the half-equation for the side-reaction occurring at the cathode. [1]

A current of 5.0 A is supplied over 2 hours during charging.

- (iii) Use the information given and your answer in (b)(i) to calculate the amount of electrons consumed by the side-reaction. [1]
- (iv) Suggest why the battery needs to be replaced after about 1000 charge-discharge cycles. [1]



(c) Chloroethanal is the starting material in the synthesis of aspartic acid.



- (i) Suggest structures for the intermediates **H** and **K**. [2]
- (ii) Suggest reagents and conditions for step 2 and for step 4. [2]
- (iii) The reaction in step 2 produces sample **J**, which does not show optical activity. Explain the observation. [2]

[Total: 18]

## Section B

Answer **one** question from this section

- 4 (a) For many compounds the enthalpy change of formation cannot be calculated directly. An indirect method based on enthalpy changes of combustion can be used.

The enthalpy change of combustion can be found by a calorimetry experiment in which the heat energy given off during combustion is used to heat a known mass of water and the temperature change recorded.

- (i) Define the term *standard enthalpy change of combustion*. [1]

- (ii) Write the equation for the complete combustion of ethanol,  $\text{CH}_3\text{CH}_2\text{OH}$ . [1]

In an experiment to determine the enthalpy change of combustion of ethanol, 0.23 g of ethanol was burned and the heat given off raised the temperature of 100 g of water by  $16.3^\circ\text{C}$ .


- (iii) Calculate the heat energy change during the combustion of ethanol. [1]

- (iv) Hence, calculate the enthalpy change on burning 1 mole of ethanol. [2]

- (v) Suggest one reason why the value for the enthalpy change of combustion of ethanol determined by a simple laboratory calorimetry experiment is likely to be lower than the true value. [1]

- (b) Epoxides are cyclic ethers commonly used in organic reactions.


Ethylene carbonate can be prepared from an epoxide, oxirane by the following reactions.



- (i) Suggest the type of reaction for reaction 2. [1]

- (ii) Reaction 1 is an acid-catalysed reaction that proceeds via a three-step mechanism:

- 1) Protonation of oxirane by  $\text{H}_3\text{O}^+$ .
- 2) Ring opening of protonated oxirane due to nucleophilic attack by  $\text{H}_2\text{O}$  to




- 3) Deprotonation of the oxonium ion to yield the product, with the regeneration of  $\text{H}_3\text{O}^+$ .

Suggest the mechanism for Reaction 1.

[3]

Compound **L** can be synthesised from an epoxide in a similar manner as ethylene carbonate.



Compound **L**

- (iii) Draw the structure of the epoxide used for synthesising compound **L**. [1]
- (iv) Suggest why compound **L** is able to exhibit cis-trans isomerism. [1]
- (c) A number of isomers with the formula  $\text{Fe}(\text{H}_2\text{O})_6\text{Cl}_3$  exist. Their general formula is  $[\text{Fe}(\text{H}_2\text{O})_{6-n}\text{Cl}_n]_m\text{Cl}_{3-n}\cdot n\text{H}_2\text{O}$ .

Each isomer contains a six co-ordinated  $\text{Fe}(\text{III})$  ion in an octahedral complex. Water molecules not directly bonded with the Fe atom are held in the crystal lattice as water of crystallisation.

- (i) Similar to organic compounds, octahedral complexes can also exhibit stereoisomerism depending on the orientation of the ligands.


One such example will be the iron complex when  $n$  is 2. It can exist in two isomeric forms where only one of them has a dipole moment.

Name the type of isomerism shown by the complex.

[1]

- (ii) Draw the structures of the two isomeric forms of the complex. [2]
- (iii) State which isomer has a dipole moment. Explain your answer. [2]

- (iv) The following diagram shows how the d-orbitals are split in an octahedral environment.



When the  $\text{H}_2\text{O}$  ligand is changed to a  $\text{Cl}^-$  ligand, the  $\text{Fe}^{2+}$  ion changes the electronic configuration from a 'high spin' to a 'low spin' state.

In a 'high spin' state, the electrons occupy all the d-orbitals singly, before starting to pair up in the lower energy d-orbitals.

In a 'low spin' state, the lower energy d-orbitals are filled first, by pairing up if necessary, before the higher energy d-orbitals are used.

Use diagrams like the one above to show the electronic distribution of a  $\text{Fe}^{2+}$  ion in a high spin state, and in a low spin state.

[2]

- (v) State and explain which ligand will result in a larger energy gap,  $E$ , between its d-orbitals.

[1]

[Total: 20]

- 5 Carbonyl compounds are common in our everyday lives. They are mainly used as solvents, perfumes and flavouring agents or as intermediates in the manufacture of plastics and pharmaceuticals.

(a) The characteristic smell of cherries and fresh almonds is due to benzaldehyde.

(i) Benzaldehyde can react with chloromethane to form 3-methylbenzaldehyde. Describe the mechanism for this reaction.

[3]

(ii) Benzene can also react under a similar reaction with chloromethane.

State and explain whether benzene or benzaldehyde would react with chloromethane more readily.

[2]

(b) Benzaldehyde can also react with hot acidified dichromate(VI) to give benzoic acid. In benzene, benzoic acid associates to form dimers.



(i) Draw a diagram to illustrate the bonding in the dimer.

[1]


(ii) Predict and explain whether the dimerisation is favoured at a high or low temperature.

[2]

(iii) Suggest why the above equilibrium cannot be established in an aqueous medium.

[1]

(c) Carbonyl compounds can be prepared from alkenes via the ozonolysis reaction as shown below.



An alkene **M**,  $\text{C}_{11}\text{H}_{14}$  was treated with  $\text{O}_3$ , followed by  $\text{Zn}$  and  $\text{H}_3\text{O}^+$  to give **N**  $\text{C}_4\text{H}_8\text{O}$  and **P**,  $\text{C}_7\text{H}_6\text{O}$ . **N** gives a yellow precipitate with aqueous alkaline iodine while **P** gives a grey precipitate with ammonical silver nitrate solution. In the presence of  $\text{OH}^-$ , **P** undergoes a reaction to give **Q**,  $\text{C}_7\text{H}_6\text{O}_2$  and **S**,  $\text{C}_7\text{H}_8\text{O}$ . Both **Q** and **S** react with  $\text{Na}$  metal, but only **Q** reacts with  $\text{NaHCO}_3$ .

Suggest the structures of **M**, **N**, **P**, **Q** and **S**.

[5]

- (d) **T, U, V** and **W** are four consecutive elements in the **fourth** period of the Periodic Table. (The letters are **not** the actual symbols of the elements.)


**T** is a soft, silvery metal with a melting point just above room temperature. Its amphoteric oxide,  $T_2O_3$ , has a melting point of 1900 °C and can be formed by heating **T** in oxygen.

**W** is a solid that can exist as several allotropes, most of which contain  $W_8$  molecules. **W** burns in air to form  $WO_2$  and  $WO_3$ , which dissolves in water to form an acidic solution. The acidic solutions react with sodium hydroxide to form the salt  $Na_2WO_3$  and  $Na_2WO_4$  respectively.

- (i) Suggest the identities of **T** and **W**. [2]
- (ii) Write equations for the reactions of  $T_2O_3$  with
- hydrochloric acid,
  - sodium hydroxide
- [2]
- (iii) Suggest the structure in  $T_2O_3$ . [1]
- (iv) Write an equation for the formation of the acidic solution when  $WO_3$  dissolves in water. [1]

[Total: 20]

**BLANK PAGE**



INNOVA JUNIOR COLLEGE  
JC 2 PRELIMINARY EXAMINATION  
in preparation for General Certificate of Education Advanced Level  
**Higher 2**

---

**CHEMISTRY**

**9729/03**

Paper 3 Free Response

**10 September 2018**

**2 hours**

Candidates answer on separate paper.

Additional Materials: Writing Papers  
*Data Booklet*  
Cover Page

---

**READ THESE INSTRUCTIONS FIRST**

Write your name and class on all the work you hand in.  
Write in dark blue or black pen on both sides of the paper.  
You may use a soft pencil for any diagrams, graphs or rough working.  
Do not use staples, paper clips, highlighters, glue or correction fluid.

**Section A**

Answer **all** questions.

**Section B**

Answer **one** question.

You are advised to show all working in calculations.  
You are reminded of the need for good English and clear presentation in your answers.  
You are reminded of the need for good handwriting.  
Your final answers should be in 3 significant figures.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in the brackets [ ] at the end of each question or part question.

---

This document consists of xx printed pages.





## Section A

Answer **all** the questions in this section.

1 This question is about the chemistry of ethene and its derivatives.

(a) Ethene is the starting material to form ethanedioic acid.

(i) Suggest the synthetic route for the formation of ethanedioic acid from ethene. [2]



Step 1: cold,  $\text{KMnO}_4$ ,  $\text{NaOH}(\text{aq})$  [✓]

Step 2:  $\text{K}_2\text{Cr}_2\text{O}_7$ , dil  $\text{H}_2\text{SO}_4$ , heat [✓]

(ii) 0.200 mol of ethene is stored in a 20.0 dm<sup>3</sup> flask with 0.800 mol of ethane at 127°C.

Calculate the total pressure in the flask. Hence or otherwise, calculate the partial pressure of ethene in the flask. [2]

$$P(20.0 \times 10^{-3}) = (0.200 + 0.800)(8.31)(127 + 273)$$

$$P = 166200 \text{ Pa} \text{ [1]}$$

$$\text{Partial pressure of ethene} = 0.2/1.0 \times 166200 = 33240$$

$$= 33.2 \times 10^3 \text{ Pa} \text{ [1]}$$

(iii) The total pressure that you calculated in (a)(ii) is different from the actual pressure exerted. Suggest an explanation for the difference. [1]

There is significant intermolecular forces of attraction present between the gases OR intermolecular forces of attraction between the gases are not negligible [1].

(iv) Ethene reacts with hydrogen in the presence of nickel catalyst to form ethane.

Explain why nickel can be used as a catalyst in this reaction. [2]

Nickel has **available 3d electrons for bond formation** with the reactant molecules (ie. ethene and hydrogen) [1] and **available low lying vacant orbitals or energetically accessible orbitals which can accept electron pairs** from the reactant molecules. [1]

(v) Outline the mode of action of the catalyst in this reaction. [2]

The catalyst is in a solid state and it functions as a heterogeneous catalyst [✓] as it is in a different phase than ethene and hydrogen. Ethene and hydrogen (reactants) will be adsorbed to the surface of the Ni catalyst. [✓] Bonds in the reactant molecules are weakened [✓] which lowers the activation energy. New bonds are then formed between adjacent reactant molecules to form ethane (product). The product formed will be desorbed from the surface of the Ni catalyst. [✓]

(b) Dissolving  $4.82 \times 10^{-5}$  mol calcium ethanedioate,  $\text{CaC}_2\text{O}_4$ , in 1 dm<sup>3</sup> of water forms a saturated solution.

(i) Write an expression for the solubility product of calcium ethanedioate and state its units. [1]

$$K_{sp}(\text{CaC}_2\text{O}_4) = [\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}] \quad \text{unit: mol}^2 \text{ dm}^{-6} \quad [1]$$

- (ii) Calculate the solubility product of calcium ethanedioate. [1]

$$\begin{aligned} K_{sp} &= (4.82 \times 10^{-5})^2 \\ &= 2.32 \times 10^{-9} \quad [1] \end{aligned}$$

[units not nec. as already stated in (i)]

- (iii) 50.0 cm<sup>3</sup> of 0.100 mol dm<sup>-3</sup> CaCl<sub>2</sub> and 50.0 cm<sup>3</sup> of 0.300 mol dm<sup>-3</sup> Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub> are mixed together. Determine if CaC<sub>2</sub>O<sub>4</sub> will be precipitated out. [2]

For precipitation to occur, IP of CaC<sub>2</sub>O<sub>4</sub> must exceed K<sub>sp</sub> of CaC<sub>2</sub>O<sub>4</sub>  
Since the volume used is the same, the new concentration is halved the original concentration.


$$\text{New } [\text{Ca}^{2+}] = 0.100 / 2 = 0.05 \quad [\checkmark]$$

$$\text{New } [\text{C}_2\text{O}_4^{2-}] = 0.300 / 2 = 0.15 \quad [\checkmark]$$


$$\text{IP of CaC}_2\text{O}_4 = (0.05)(0.15) = 7.50 \times 10^{-3} \quad [\checkmark]$$

Since IP > K<sub>sp</sub>, CaC<sub>2</sub>O<sub>4</sub> will ppt out. [✓][ECF with student's K<sub>sp</sub> value from (ii)]


- (c) Synthesis of the unionised form of EDTA (H<sub>4</sub>Y) can be carried out in the laboratory using ethene as the starting material.



- (i) Draw the structures of **A** and **B**. [2]




4



- (ii) Suggest appropriate reagents and conditions for step 3.


Step 3: dilute  $\text{H}_2\text{SO}_4$ ,  $\text{KMnO}_4(\text{aq})/\text{K}_2\text{Cr}_2\text{O}_7(\text{aq})$ , heat under reflux [1] (followed by careful addition of  $\text{NaOH}(\text{aq})$ ) [1]

- (d) Ethene can be used to form ethanal. Ethanal in turn is used to synthesize glycolic acid via the reaction shown below.



- (i) Identify the intermediate **C** and state the reagent and condition for Step 3.

Step 3:  $\text{NaOH}(\text{aq})$ , heat under reflux; followed by aq.  $\text{HCl}$  [1]




- (ii) Explain the difference in acidity between glycolic acid and ethanoic acid.

In glycolic acid, the electron withdrawing  $-\text{OH}$  [✓] group reduces the negative charge on  $-\text{COO}^-$  [✓] of the conjugate base of glycolic acid, thus making the anion of glycolic acid is more stable than the ethanoate anion [✓]. Hence, glycolic acid is a stronger acid [✓]. [2]

- (iii) Two molecules of glycolic acid can react with one another under suitable conditions to form a neutral compound with the loss of two water molecules.

Suggest a possible structure for the compound formed.



- (iv) Suggest a simple chemical test to distinguish between ethanoic acid and glycolic acid.

[2]

Test:  $K_2Cr_2O_7$ , dil  $H_2SO_4$ , heat [1]

Ethanoic acid: orange  $K_2Cr_2O_7$  turns green [✓]

Glycolic acid: orange  $K_2Cr_2O_7$  remains [✓]

OR

Test:  $KMnO_4$ , dil  $H_2SO_4$ , heat

Ethanoic acid: purple  $KMnO_4$  remains

Glycolic acid: purple  $KMnO_4$  decolorised. Effervescence observed. gas produced forms white ppt with  $Ca(OH)_2$


[Total: 23]

- 2 Iron is the fourth most common element in the Earth's crust. It is a d-block element which is known to exhibit different characteristics from the s-block elements. Since ancient times, iron has been widely employed in a variety of applications.


- (a) (i) One well-known property of iron and its compounds is the ability to catalyse reactions. For example, aqueous iron(II) chloride can be used to catalyse the reaction between  $I^-$  and  $S_2O_8^{2-}$ , to form  $I_2$  and  $SO_4^{2-}$  ions.

Using relevant  $E^\ominus$  values from the *Data Booklet*, explain why iron(II) chloride can be used as a catalyst for this reaction.

[2]



$$\begin{aligned} E_{\text{cell}} &= +2.01 - (0.77) \\ &= +1.24 \text{ V (reaction is energetically feasible) [1]} \end{aligned}$$



$$\begin{aligned} E_{\text{cell}} &= +0.77 - (+0.54) \\ &= +0.23 \text{ V (reaction is energetically feasible) [1]} \end{aligned}$$

- (ii) A sample of iron was vapourised, ionised and passed through an electric field. Analysis of deflection occurring at the electric field region revealed that a sample of  $^{32}S^{2-}$  ions would be deflected by  $+20^\circ$  towards the positive potential.

What is the angle, and direction of deflection for a sample of  $^{56}Fe^{3+}$  ions passing through the same electric field?

[2]

Angle of deflection of  $^{32}S^{2-} = 20$

$$20 = k (-2 / 32)$$


$$k = -320$$

Angle of deflection of  $^{56}Fe^{3+} = (-320) (3 / 56)$

$$= -17.1^\circ \quad [1 \text{ for final answer; 1m for indication of direction}]$$

or  $17.1^\circ$  [1m] towards the negative potential [1m]

- (b) Ferrocene,  $\text{Fe}(\text{C}_5\text{H}_5)_2$ , is an orange solid which is known to exhibit anti-cancer activity. In this complex,  $\text{C}_5\text{H}_5^-$  is the ligand and it donates  $\pi$  electrons from the ring to the vacant 3d orbital of Fe. The structure of ferrocene is given below.



- (i) State the oxidation number of Fe in ferrocene. [1]  
+2

- (ii) Suggest why ferrocene is a coloured complex. [2]  
There is partially filled 3d orbitals in  $\text{Fe}^{2+}$ . [✓]

In the presence of ligands, 3d orbitals of iron are split into 2 groups with small energy gap (d–d splitting). [✓]

Some light energy is used to promote an electron [✓] from a d-orbital of lower energy to a d-orbital of a higher energy [✓].

(Colour observed is complementary to the colour absorbed.)

- (iii) Light of a longer wavelength is lower in energy than light of a shorter wavelength. The following table shows the spectral colours and the corresponding wavelengths.


Colour	Wavelength / nm
Violet	380 – 450
Blue	450 – 495
Green	495 – 570
Yellow	570 – 590
Orange	590 – 620
Red	620 – 750

Given that aqueous  $\text{Fe}^{2+}$  ion is green in colour, suggest and explain if water causes a larger split between the two groups of 3d orbitals as compared to  $\text{C}_5\text{H}_5^-$ . [2]

Water causes a smaller split between the 3d orbitals. [1]

With water as the ligand, electromagnetic waves with energy corresponding to red is absorbed which has a lower energy than blue, which is absorbed when  $\text{C}_5\text{H}_5^-$  is the ligand. [1]


- (c) Ethylbenzene is used to synthesize (3-chlorophenyl)methanol which is used as a general solvent for inks, paints, lacquers, epoxy resin coatings and as a degreasing agent.



Starting with ethylbenzene, outline a three-step reaction scheme to obtain (3-chlorophenyl)methanol. Your answer should include clearly the reagents and conditions in each step, and the structures of all intermediates formed.

[3]

Ans:



5 [✓] – 3m  
 3 to 4 [✓] – 2m  
 2 [✓] – 1m


- (d) **D** is an achiral organic compound with the molecular formula  $\text{C}_2\text{H}_7\text{NO}$ . It can be formed from the reaction between a primary amide and lithium aluminium hydride. In the presence of a suitable catalyst, 1 mole of **D** reacts with 1 mole of benzoic acid to form **E**,  $\text{C}_9\text{H}_{11}\text{NO}_2$ . However, 1 mole of **D** requires 2 moles of benzoyl chloride to react completely to form **F**,  $\text{C}_{16}\text{H}_{15}\text{NO}_3$  and copious white fumes. 2 moles of **D** can also react with gaseous  $\text{PCl}_5$  to form a cyclic **G**,  $\text{C}_4\text{H}_{10}\text{N}_2$ , which contains a 6-membered ring.

Deduce the structures of **D**, **E**, **F** and **G** and explain the reactions described.

[7]


Info	Deduction
<b>D</b> is formed from reaction between primary amide and LiAlH <sub>4</sub>	<b>D</b> contains a <u>primary amine</u> [✓]
1 mol <b>D</b> reacts with 1 mol of benzoic acid to form <b>E</b> , C <sub>9</sub> H <sub>11</sub> NO <sub>2</sub>	<ul style="list-style-type: none"> <li><b>D</b> undergoes <u>condensation</u> / nucleophilic acyl substitution [✓]</li> <li><b>E</b> is an <u>ester</u> [✓]</li> <li><b>D</b> contains <u>alcohol</u> [✓]</li> </ul>
1 mol <b>D</b> require 2 mol benzoyl chloride to react completely to form Compound <b>F</b> , C <sub>16</sub> H <sub>15</sub> NO <sub>3</sub> and copious white fumes.	<ul style="list-style-type: none"> <li><b>D</b> undergoes <u>condensation</u>/ nucleophilic acyl substitution [✓]</li> <li>Copious white fumes are <u>HCl</u> [✓]</li> <li><b>F</b> contains an <u>amide</u> [✓] and an <u>ester</u> [✓]</li> </ul>
2 mol <b>D</b> reacts with gaseous PCl <sub>5</sub> to form a cyclic Compound <b>G</b> , C <sub>4</sub> H <sub>10</sub> N <sub>2</sub>	<ul style="list-style-type: none"> <li><b>D</b> undergoes <u>nucleophilic substitution</u> [✓]</li> <li><b>G</b> contains an <u>amine</u> [✓]</li> </ul>
8 - 10 [✓] – 3m 5 - 7 [✓] – 2m 2 - 4 [✓] – 1m	

**D** (C<sub>2</sub>H<sub>7</sub>NO) : NH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH [1]



Compound **E** (C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>) :


[1]



Compound **F** (C<sub>16</sub>H<sub>15</sub>NO<sub>3</sub>) :

[1]

ECF for structure E and F based on wrong structure D, H<sub>2</sub>NCH(OH)CH<sub>3</sub>.





Compound **G** (C<sub>4</sub>H<sub>10</sub>N<sub>2</sub>) :


[1]

[Total: 19]

- 3 An organic radical battery (ORB) is a relatively new type of battery which uses flexible plastics, to provide electrical power. One type of hybrid ORB/Li-ion battery consists of: a cathode made from solid organic polymer containing oxoammonium ion formed from (2,2,6,6-Tetramethylpiperidin-1-yl)oxyl (TEMPO) nitroxide radicals; an anode made from graphite with Li atoms inserted between the graphite layers; and an electrolyte of  $\text{LiPF}_6$  dissolved in organic solvent.

During discharge, Li atoms give up electrons at the anode to become  $\text{Li}^+$  ions. The electrons travel round the external circuit, and are picked up by the cathode. The anions and cations in the electrolyte move to the anode and cathode respectively. This is illustrated in the following

diagram in which  and  are simplified representations of the polymer containing oxoammonium ion and TEMPO nitroxide radicals respectively.



- (a) (i) Graphite is often mixed in the polymer used for making the cathode electrode.

Suggest a reason for this.

[1]



To increase the electrical conductivity of the electrode  
Or (good) conductor of electricity [1]

- (ii) Calculate the oxidation number of N in the cathode **before** discharge. [1]


Oxidation number of N is +1 [1]

- (iii) The  $E_{\text{cell}}$  generated by the hybrid ORB/Li-ion battery under standard conditions is 2.17 V.

Use relevant  $E^\ominus$  value from the *Data Booklet* to calculate the electrode potential generated by the cathode half-cell. [1]

$$\begin{aligned} E_{\text{cell}} &= E_{\text{red}} - E_{\text{ox}} \\ +2.17 &= E_{\text{red}} - (-3.04) \\ &= -0.87 \text{ V} \quad [1, \text{ with correct working and sign}] \end{aligned}$$

- (iv) During discharge, the following reaction occurs at the cathode.




where R represents the organic polymer cathode.

Write an equation for the overall process that occurs during discharge. [1]



- (v) Draw the dot-and-cross diagram of the  $\text{PF}_6^-$  ion and state its shape. [2]



[1, check for dative bond and overall charge]

Shape: Octahedral [1]

- (vi) Suggest whether  $\text{LiPF}_6$  or  $\text{LiF}$  has a lower melting point. Explain your answer. [2]

$\text{LiPF}_6$ . [1, conditional provided student considered LE)

$\text{PF}_6^-$  has a **larger anionic radius** [✓] than  $\text{F}^-$ , giving rise to a **lower magnitude**

of LE [✓] (since L.E.  $\propto \left| \frac{q_+ \times q_-}{r_+ + r_-} \right|$ ). Hence,  $\text{LiPF}_6$  has a lower ionic bond strength

and less energy is required to overcome the ionic bond, giving rise to lower mp.

- (b) The hybrid ORB/Li-ion battery is a secondary battery, i.e., it is rechargeable.

During charging, 1.22 g of Li is regenerated from  $\text{Li}^+$  ions at the cathode.

- (i) Calculate the amount of electrons required to form 1.22 g of Li. [1]

$$\text{Amount of Li formed} = 1.22 / 6.9 = 0.1768 \text{ mol}$$




$$\text{Amount of electrons consumed} = 0.1768 = 0.177 \text{ mol [1].}$$

Besides the generation of Li, there is a competing side-reaction that occurs at the cathode.

In this side-reaction, ethylene carbonate,  $\text{C}_3\text{H}_4\text{O}_3$  undergoes reduction in the presence of  $\text{Li}^+$  ions to form ethene and lithium carbonate.

- (ii) Write the half-equation for the side-reaction occurring at the cathode. [1]



A current of 5.0 A is supplied over 2 hours during charging.

- (iii) Use the information given and your answer in (b)(i) to calculate the amount of electrons consumed by the side-reaction. [1]


$$\text{Amount of electrons supplied} = (5.0) (2 \times 3600) / 96500 = 0.3731 \text{ mol [✓].}$$

$$\begin{aligned} \text{Hence, amount of electrons consumed by side-reaction} \\ &= 0.3731 - 0.1768 \\ &= 0.196 \text{ mol [✓, allow ecf from (i)].} \end{aligned}$$

- (iv) Suggest why the battery needs to be replaced after about 1000 charge-discharge cycles. [1]

Some of the Li ions are depleted / not all the Li metal is regenerated during charging due to the side-reaction [1].

- (c) Chloroethanal is the starting material in the synthesis of aspartic acid.



- (i) Suggest structures for the intermediates **H** and **K**. [2]



- (ii) Suggest reagents and conditions for step 2 and for step 4. [2]

Step 2: HCN, trace NaOH(aq)/NaCN, cold [1]

Step 4: excess, concentrated ethanolic NH<sub>3</sub>, heat in sealed tube [1]

- (iii) The reaction in step 2 produces sample **J**, which does not show optical activity. Explain the observation. [2]

The nucleophile, CN<sup>-</sup>, has equal chances of attacking either above or below the plane of the electron-deficient carbonyl carbon in compound H [1]. Thus, the product mixture contains 50% of each of the enantiomer, forming a racemic mixture [1]. Hence, the optical activity of the 2 enantiomers cancels out [1]. Thus, sample **J** produced does not show optical activity.

[Total: 18]

### Section B

Answer **one** question from this section


- 4 (a) For many compounds the enthalpy change of formation cannot be calculated directly. An indirect method based on enthalpy changes of combustion can be used.

The enthalpy change of combustion can be found by a calorimetry experiment in which the heat energy given off during combustion is used to heat a known mass of water and the temperature change recorded.

- (i) Define the term *standard enthalpy change of combustion*. [1]

Standard enthalpy change of combustion,  $\Delta H_c^\ominus$ , of a compound is the enthalpy change (heat evolved) when 1 mole of the substance is completely burnt in oxygen under standard conditions of 298K and 1 bar. [1]

- (ii) Write the equation for the complete combustion of ethanol, CH<sub>3</sub>CH<sub>2</sub>OH. [1]



In an experiment to determine the enthalpy change of combustion of ethanol, 0.23 g of ethanol was burned and the heat given off raised the temperature of 100 g of water by 16.3 °C.

- (iii) Calculate the heat energy change during the combustion of ethanol. [1]

$$\begin{aligned} \text{Heat change} &= mc\Delta T \\ &= (100)(4.18)(16.3) \\ &= 6813.4 \end{aligned}$$

= 6810 J[1]  
(no mark awarded if unit given is wrong)

- (iv) Hence, calculate the enthalpy change on burning 1 mole of ethanol. [2]

Amount of ethanol =  $0.23 / (2 \times 12.0 + 6.0 + 16.0)$   
=  $5.00 \times 10^{-3}$  mol


Enthalpy change =  $- (6813.4 / 5.00 \times 10^{-3})$  [1m; ECF from (iii)]  
=  $- 1362680 \text{ J mol}^{-1}$   
=  $- 1360 \text{ kJ mol}^{-1}$  [1m include unit & sign]

- (v) Suggest one reason why the value for the enthalpy change of combustion of ethanol determined by a simple laboratory calorimetry experiment is likely to be lower than the true value. [1]

Heat loss to the surrounding. / Incomplete combustion/ Ethanol is volatile and evaporated./ Ethanol is impure. / Not all energy is absorbed by the water.  
(Any answer 1 mark)

- (b) Epoxides are cyclic ethers commonly used in organic reactions.

Ethylene carbonate can be prepared from an epoxide, oxirane by the following reactions.




- (i) Suggest the type of reaction for reaction 2. [1]  
Condensation/ nucleophilic acyl substitution [1]
- (ii) Reaction 1 is an acid-catalysed reaction that proceeds via a three-step mechanism:

- 1) Protonation of oxirane by  $\text{H}_3\text{O}^+$ .
- 2) Ring opening of protonated oxirane due to nucleophilic attack by  $\text{H}_2\text{O}$  to yield an oxonium ion, OCC[OH2+].
- 3) Deprotonation of the oxonium ion to yield the product, with the regeneration of  $\text{H}_3\text{O}^+$ .

Suggest the mechanism for Reaction 1.

[3]




- ✓ - correct half-arrows and lone pairs for step 1
- ✓ - formation of H<sub>2</sub>O in step 1
- ✓ - correct structure for protonated epoxide in step 1
- ✓ - correct half-arrows and lone pair for step 2
- ✓ - correct half-arrows and regeneration of H<sub>3</sub>O<sup>+</sup> for step 3

(fast & slow labels not marked for)


3m – all 5 ✓ ; 2m – 3 to 4 ✓ ; 1m – 2 ✓

Compound **L** can be synthesised from an epoxide in a similar manner as ethylene carbonate.



Compound **L**

- (iii) Draw the structure of the epoxide used for synthesising compound **L**. [1]



- (iv) Suggest why compound **L** is able to exhibit cis-trans isomerism. [1]  
**Restricted C-C bond rotation [1] due to ring strain/ rigidity of ring**  
 (and both C atoms of the ring is attached to 2 different groups or atoms)

- (c) A number of isomers with the formula Fe(H<sub>2</sub>O)<sub>6</sub>Cl<sub>3</sub> exist. Their general formula is [Fe(H<sub>2</sub>O)<sub>6-n</sub>Cl<sub>n</sub>](Cl<sub>3-n</sub>).nH<sub>2</sub>O.

Each isomer contains a six co-ordinated Fe(III) ion in an octahedral complex. Water molecules not directly bonded with the Fe atom are held in the crystal lattice as water of crystallisation.

- (i) Similar to organic compounds, octahedral complexes can also exhibit stereoisomerism depending on the orientation of the ligands.


One such example will be the iron complex when  $n$  is 2. It can exist in two isomeric forms where only one of them has a dipole moment.

Name the type of isomerism shown by the complex

[1]

Cis- Trans isomerism

- (ii) Draw the structures of the two isomeric forms of the complex. [2]




[1 mark each] (labelling of cis-trans not required)

- (iii) State which isomer has a dipole moment. Explain your answer. [2]

Cis isomer has a dipole moment. [1] (student need not mention "cis". Marks are awarded as long as student identify the correct isomer)  
Cl<sup>-</sup> on the same side of the complex/ overall dipole moment does not cancel out.[1]

- (iv) The following diagram shows how the d-orbitals are split in an octahedral environment.



When the  $\text{H}_2\text{O}$  ligand is changed to a  $\text{Cl}$  ligand, the  $\text{Fe}^{2+}$  ion changes the electronic configuration from a 'high spin' to a 'low spin' state.


In a 'high spin' state, the electrons occupy all the d-orbitals singly, before starting to pair up in the lower energy d-orbitals.

In a 'low spin' state, the lower energy d-orbitals are filled first, by pairing up if necessary, before the higher energy d-orbitals are used.


Use diagrams like the one above to show the electronic distribution of a  $\text{Fe}^{2+}$  ion in a high spin state, and in a low spin state.

$\text{Fe}^{2+} : 3d^6$

'High spin' state



'Low spin' state



- (v) State and explain which ligand will result in a larger energy gap,  $E$ , between its d-orbitals.

[1]

(Complex with) Cl ligand [1]

Electrons will only pair up in the lower orbitals when the energy gap was greater than the interelectronic repulsion between the two electrons in the same orbital. (wtse) [1]

[Total: 20]



- 5 Carbonyl compounds are common in our everyday lives. They are mainly used as solvents, perfumes and flavouring agents or as intermediates in the manufacture of plastics and pharmaceuticals.

(a) The characteristic smell of cherries and fresh almonds is due to benzaldehyde.

- (i) Benzaldehyde can react with chloromethane to form 3-methylbenzaldehyde. Describe the mechanism for this reaction.

[3]

Electrophilic Substitution [1]



- (ii) Benzene can also react under a similar reaction with chloromethane.

State and explain whether benzene or benzaldehyde would react with chloromethane more readily.

[2]

Benzene would react **more readily** [✓] with chloromethane.


Benzaldehyde contains an **electron withdrawing CHO group** [✓] which **decreases the electron density of the benzene ring** [✓], making it **less susceptible to electrophilic attacks** [✓].

- (b) Benzaldehyde can also react with hot acidified dichromate(VI) to give benzoic acid. In benzene, benzoic acid associates to form dimers.



- (i) Draw a diagram to illustrate the bonding in the dimer.

[1]



All details included (dipoles, lone pair, hydrogen bonds) – [1]

- (ii) Predict and explain whether the dimerisation is favoured at a high or low temperature.

[2]


**Low temperature [1, conditional].** At low temperature, the position of equilibrium shifts to the right to favour the **exothermic reaction [1]** to release heat.

- (iii) Suggest why the above equilibrium cannot be established in an aqueous medium.

In aqueous medium, **benzoic acid would form hydrogen bonds with the water molecules instead** of forming a dimer, [1]

[1]

- (c) Carbonyl compounds can be prepared from alkenes via the ozonolysis reaction as shown below.




An alkene **M**,  $\text{C}_{11}\text{H}_{14}$  was treated with  $\text{O}_3$ , followed by  $\text{Zn}$  and  $\text{H}_3\text{O}^+$  to give **N**  $\text{C}_4\text{H}_8\text{O}$  and **P**,  $\text{C}_7\text{H}_6\text{O}$ . **N** gives a yellow precipitate with aqueous alkaline iodine while **P** gives a grey precipitate with ammonical silver nitrate solution. In the presence of  $\text{OH}^-$ , **P** undergoes a reaction to give **Q**,  $\text{C}_7\text{H}_6\text{O}_2$  and **S**,  $\text{C}_7\text{H}_8\text{O}$ . Both **Q** and **S** react with  $\text{Na}$  metal, but only **Q** reacts with  $\text{NaHCO}_3$ .

Suggest the structures of **M**, **N**, **P**, **Q** and **S**.

[5]





[1] each

- (d) **T, U, V** and **W** are four consecutive elements in the **fourth** period of the Periodic Table. (The letters are **not** the actual symbols of the elements.)



**T** is a soft, silvery metal with a melting point just above room temperature. Its amphoteric oxide,  $T_2O_3$ , has a melting point of  $1900\text{ }^\circ\text{C}$  and can be formed by heating **T** in oxygen.

**W** is a solid that can exist as several allotropes, most of which contain  $W_8$  molecules. **W** burns in air to form  $WO_2$  and  $WO_3$ , which dissolves in water to form an acidic solution. The acidic solutions react with sodium hydroxide to form the salt  $Na_2WO_3$  and  $Na_2WO_4$  respectively.

- (i) Suggest the identities of **T** and **W**. [2]

**T = Ga [1]**      **W = Se [1]**

- (ii) Write equations for the reactions of  $T_2O_3$  with
- hydrochloric acid,
  - sodium hydroxide
- [2]




(in writing the equations, students can replace **T** with Ga)

- (iii) Suggest the structure in  $T_2O_3$ . [1]

Giant Ionic Structure [1]

- (iv) Write an equation for the formation of the acidic solution when  $WO_3$  dissolves in water. [1]



(in writing the equation, students can replace **W** with Se)

[Total: 20]